

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number  
**WO 02/088097 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 249/06**,  
249/08, 249/10, 403/04, 401/14, 405/14, 417/14, 403/14

(21) International Application Number: PCT/US02/13396

(22) International Filing Date: 26 April 2002 (26.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/287,204 27 April 2001 (27.04.2001) US

(71) Applicant (for all designated States except US): **VERTEX  
PHARMACEUTICALS INCORPORATED** [US/US];  
130 Waverly Street, Cambridge, MA 02139-4242 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HALE, Michael,  
Robin** [US/US]; 42 Sunset Road, Bedford, MA 01730  
(US). **MALTAIS, Francois** [CA/US]; 24 Randolph Drive,  
Tewksbury, MA 01876 (US).

(74) Agents: **HALEY, James, F.** et al.; c/o Fish & Neave, 1251  
Avenue of the Americas, New York, NY 10020 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

**Published:**

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TRIAZOLE-DERIVED KINASE INHIBITORS AND USES THEREOF

(57) Abstract: Described herein are compounds that are useful as protein kinase inhibitors having the formula: wherein Ht, R<sup>2</sup>, T, and m are as described in the specification. The compounds are useful for treating diseases in mammals that are alleviated by a protein kinase inhibitor, particularly diseases such as cancer, inflammatory disorders, restenosis, and cardiovascular disease.



WO 02/088097 A1

TRIAZOLE-DERIVED KINASE INHIBITORS AND USES THEREOFCROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to United States Provisional Application 60/287,204, filed April 27, 2001, the contents of which are incorporated herein by reference.

TECHNICAL FIELD OF THE INVENTION

5           The present invention is in the field of medicinal chemistry and relates to triazole compounds that are protein kinase inhibitors, especially inhibitors of ERK, compositions containing such compounds and methods of use. The compounds are useful for treating  
10 cancer and other diseases that are alleviated by protein kinase inhibitors.

BACKGROUND OF THE INVENTION

Mammalian mitogen-activated protein (MAP)  
15 kinases are serine/threonine kinases that mediate intracellular signal transduction pathways (Cobb and Goldsmith, 1995, *J Biol. Chem.*, **270**, 14843; Davis, 1995, *Mol. Reprod. Dev.* **42**, 459). Members of the MAP kinase family share sequence similarity and conserved structural  
20 domains, and include the ERK2 (extracellular signal regulated kinase), JNK (Jun N-terminal kinase), and p38 kinases. JNKs and p38 kinases are activated in response to the pro-inflammatory cytokines TNF-alpha and interleukin-1, and by cellular stress such as heat shock,  
25 hyperosmolarity, ultraviolet radiation, lipopolysaccharides and inhibitors of protein synthesis (Derijard et al., 1994, *Cell* **76**, 1025; Han et al., 1994,

*Science* **265**, 808; Raingeaud et al., 1995, *J Biol. Chem.* **270**, 7420; Shapiro and Dinarello, 1995, *Proc. Natl. Acad. Sci. USA* **92**, 12230). In contrast, ERKs are activated by mitogens and growth factors (Bokemeyer et al. 1996, *Kidney Int.* **49**, 1187).

ERK2 is a widely distributed protein kinase that achieves maximum activity when both Thr183 and Tyr185 are phosphorylated by the upstream MAP kinase kinase, MEK1 (Anderson et al., 1990, *Nature* **343**, 651; Crews et al., 1992, *Science* **258**, 478). Upon activation, ERK2 phosphorylates many regulatory proteins, including the protein kinases Rsk90 (Bjorbaek et al., 1995, *J. Biol. Chem.* **270**, 18848) and MAPKAP2 (Rouse et al., 1994, *Cell* **78**, 1027), and transcription factors such as ATF2 (Raingeaud et al., 1996, *Mol. Cell Biol.* **16**, 1247), Elk-1 (Raingeaud et al. 1996), c-Fos (Chen et al., 1993 *Proc. Natl. Acad. Sci. USA* **90**, 10952), and c-Myc (Oliver et al., 1995, *Proc. Soc. Exp. Biol. Med.* **210**, 162). ERK2 is also a downstream target of the Ras/Raf dependent pathways (Moodie et al., 1993, *Science* **260**, 1658) and may help relay the signals from these potentially oncogenic proteins. ERK2 has been shown to play a role in the negative growth control of breast cancer cells (Frey and Mulder, 1997, *Cancer Res.* **57**, 628) and hyperexpression of ERK2 in human breast cancer has been reported (Sivaraman et al., 1997, *J Clin. Invest.* **99**, 1478). Activated ERK2 has also been implicated in the proliferation of endothelin-stimulated airway smooth muscle cells, suggesting a role for this kinase in asthma (Whelchel et al., 1997, *Am. J. Respir. Cell Mol. Biol.* **16**, 589).

A number of compounds have been developed that purport to specifically inhibit various MAPKs. PCT publication WO 95/31451 describes pyrazole derivatives that inhibit p38. However, it is not clear whether these

compounds have the appropriate pharmacological profiles to be therapeutically useful.

Aryl-substituted pyrroles are known in the literature. In particular, tri-aryl pyrroles (US 5,837,719) have been described as having glucagon antagonist activity.

Heterocycle-substituted triazole compounds are known in the literature. In particular, bis([1,2,3]triazoles) (Abbasoglu et al., 1999, *Indian J. Chem., Sect. B* **38B**, 413; Klaus, 1989, *Chem. Ber.* **122**, 1175; Samsonov et al., 1993; *Khim. Geterotsikl. Soedin.* **29**, 1169) and tetrazolyl-triazole (Ried and Laoutidis, 1990, *Chem.-Ztg.* **114**, 246; Vereshchagin et al., 1984, *Zh. Org. Khim.* **20**, 142) compounds have been described.

There is a high unmet medical need to develop new therapeutic treatments that are useful in treating the various conditions associated with ERK2 activation. For many of these conditions the currently available treatment options are inadequate.

Accordingly, there is great interest in new and effective inhibitors of protein kinase, including ERK2 inhibitors, which are useful in treating various conditions associated with protein kinase activation.


#### SUMMARY OF THE INVENTION

The present invention provides compounds and compositions thereof that are useful as protein kinase inhibitors, especially as inhibitors of ERK. These compounds can be used alone or in combination with other therapeutic or prophylactic agents, such as antibiotics, immunomodulators or other anti-inflammatory agents, for the treatment or prophylaxis of diseases mediated by protein kinases, including ERK2. According to a preferred embodiment, the compounds of this invention are



capable of binding to the active site of ERK2 and inhibiting the activity of that enzyme.

It is a principal object of this invention to provide compounds that are protein kinase inhibitors  
 5 represented by formula I:



or a pharmaceutically acceptable derivative thereof, wherein:

Ht is a heterocyclic ring selected from pyrrol-3-yl, [1,2,4]triazol-3-yl, [1,2,3]triazol-4-yl, or tetrazol-5-yl, said pyrrol-3-yl having  $R^3$  and  $Q_n-R^4$  substituents, and said [1,2,4]triazol-3-yl or  
 10 [1,2,3]triazol-4-yl substituted by either  $R^3$  or  $Q_n-R^4$ ;

T and Q are each independently selected from  $-C(R^7)_2-$ ,  $-C(O)-$ ,  $-C(O)C(O)-$ ,  $-C(O)NR^7-$ ,  $-C(O)NR^7NR^7-$ ,  $-CO_2-$ ,  $-OC(O)-$ ,  $-NR^7CO_2-$ ,  $-O-$ ,  $-NR^7C(O)NR^7-$ ,  $-OC(O)NR^7-$ ,  $-NR^7NR^7-$ ,  $-NR^7C(O)-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^7-$ ,  $-SO_2NR^7-$ , or  
 15  $-NR^7SO_2-$ ,  $-NR^7SO_2NR^7-$ ;

m and n are each independently selected from zero or one;

$R^2$  is selected from hydrogen, CN, halogen, R,  $N(R^7)_2$ , OR, or OH;

20 each  $R^3$  is independently selected from  $R^7$ , F, Cl,  $-(CH_2)_yN(R^7)_2$ ,  $-N(R^7)_2$ ,  $-OR^7$ ,  $-SR^7$ ,  $-NR^7C(O)R^7$ ,  $-NR^7C(O)N(R^7)_2$ ,  $-C(O)N(R^7)_2$ ,  $-SO_2R^7$ ,  $-NR^7SO_2R^7$ ,  $-C(O)R^7$ ,  $-C(O)OR^7$ , CN or  $-SO_2N(R^7)_2$ ;

$R^4$  is selected from  $-(CH_2)_yR^6$ ,  $-(CH_2)_yCH(R^6)_2$ ,  
 25  $-(CH_2)_yCH(R^8)CH(R^6)_2$ ,  $-N(R^5)_2$ ,  $-NR^5(CH_2)_yN(R^5)_2$ ;

each R is independently selected from an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, heteroaryl having 5-10 ring atoms, and heterocyclyl having 3-10 ring atoms;

5 each R<sup>5</sup> is independently selected from R, -R<sup>7</sup>, -C(O)R<sup>7</sup>, -CO<sub>2</sub>R, -C(O)N(R<sup>7</sup>)<sub>2</sub>, -SO<sub>2</sub>R<sup>7</sup>, -(CH<sub>2</sub>)<sub>y</sub>R<sup>6</sup>, or -(CH<sub>2</sub>)<sub>y</sub>CH(R<sup>6</sup>)<sub>2</sub>;

y is 0-6;

10 each R<sup>6</sup> is independently selected from R, -(CH<sub>2</sub>)<sub>y</sub>R, -OR, -CO<sub>2</sub>R, -(CH<sub>2</sub>)<sub>y</sub>N(R<sup>7</sup>)<sub>2</sub>, -N(R<sup>7</sup>)<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -NR<sup>7</sup>C(O)N(R<sup>7</sup>)<sub>2</sub>, -C(O)N(R<sup>7</sup>)<sub>2</sub>, -SO<sub>2</sub>R<sup>7</sup>, -NR<sup>7</sup>SO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -CN, -SO<sub>2</sub>N(R<sup>7</sup>)<sub>2</sub>;

15 each R<sup>7</sup> is independently selected from hydrogen or an optionally substituted C<sub>1-6</sub> aliphatic group, or two R<sup>7</sup> on the same nitrogen are taken together with the nitrogen to form a 5-8 membered ring heterocyclyl or heteroaryl ring;

each R<sup>8</sup> is independently selected from R, -(CH<sub>2</sub>)<sub>w</sub>OR<sup>7</sup>, -(CH<sub>2</sub>)<sub>w</sub>N(R<sup>5</sup>)<sub>2</sub>, or -(CH<sub>2</sub>)<sub>w</sub>SR<sup>7</sup>;

20 each w is independently 0-4;

provided that when Ht is tetrazol-5-yl, and m is 0, R<sup>2</sup> is other than H or an unsubstituted phenyl; when m is 0 and R<sup>2</sup>, R<sup>3</sup> and Q<sub>n</sub>-R<sup>4</sup> are all H, Ht is other than [1,2,3]triazole; and when T<sub>m</sub>-R<sup>2</sup> and Q<sub>n</sub>-R<sup>4</sup> are both  
25 C(O)OH, Ht is other than [1,2,3]triazol-5-yl.

It is a further objective of this invention to provide pharmaceutical compositions comprising the protein kinase inhibitors of this invention. In a  
30 preferred embodiment, the protein kinase inhibitors inhibit ERK2. These compositions may be utilized in methods for treating or preventing a variety of protein kinase-mediated disorders, such as cancer, stroke, diabetes, hepatomegaly, cardiovascular disease including

cardiomegaly, Alzheimer's disease, cystic fibrosis, viral disease, autoimmune diseases, atherosclerosis, restenosis, psoriasis, allergic disorders including asthma, inflammation, neurological disorders and hormone-related diseases. Each of the above-described methods is also part of the present invention.

It is a further objective of this invention to provide methods for making the compounds and compositions of this invention.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to compounds of formula I. Accordingly, it has now been found that compounds of this invention and compositions thereof are effective as protein kinase inhibitors, especially as inhibitors of ERK2.

As used herein, the following definitions shall apply unless otherwise indicated. The phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and each substitution is independent of the other. Also, combinations of substituents or variables are permissible only if such combinations result in a chemically stable arrangement.

The term "chemically stable arrangement" or "chemically feasible and stable" as used herein, refers to a compound structure that renders the compound sufficiently stable to allow manufacture and administration to a mammal by methods known in the art. Typically, such compounds are stable at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

The term "aliphatic" or "aliphatic group" as used herein means a straight-chain or branched C<sub>1</sub>-C<sub>12</sub> hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a  
5 monocyclic C<sub>3</sub>-C<sub>8</sub> hydrocarbon or bicyclic C<sub>8</sub>-C<sub>12</sub> hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" or "cycloalkyl"), that has a single point of attachment to the rest of the  
10 molecule wherein any individual ring in said bicyclic ring system has 3-7 members. For example, suitable aliphatic groups include, but are not limited to, linear or branched or alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or  
15 (cycloalkyl)alkenyl.

The terms "alkyl", "alkoxy", "hydroxyalkyl", "alkoxyalkyl", and "alkoxycarbonyl", used alone or as part of a larger moiety includes both straight and branched chains containing one to twelve carbon atoms.  
20 The terms "alkenyl" and "alkynyl" used alone or as part of a larger moiety shall include both straight and branched chains containing two to twelve carbon atoms.

The terms "haloalkyl", "haloalkenyl" and "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case  
25 may be, substituted with one or more halogen atoms. The term "halogen" means F, Cl, Br, or I.

The term "heteroatom" means nitrogen, oxygen, or sulfur and includes any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen.  
30 Also the term "nitrogen" includes a substitutable nitrogen of a heterocyclic ring. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2H-pyrrolyl), NH (as

in pyrrolidinyl) or  $\text{NR}^+$  (as in N-substituted pyrrolidinyl).

The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy," or "aryloxyalkyl," refers to monocyclic, bicyclic and tricyclic carbocyclic ring systems having a total of five to fourteen members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term "aryl" may be used interchangeably with the term "aryl ring".

The term "heterocycle", "heterocyclyl", or "heterocyclic" as used herein means non-aromatic, monocyclic, bicyclic or tricyclic ring systems having five to fourteen ring members in which one or more ring members is a heteroatom, wherein each ring in the system contains 3 to 8 ring members.

The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic and tricyclic ring systems having a total of five to fourteen ring members, and wherein: 1) at least one ring in the system is aromatic; 2) at least one ring in the system contains one or more heteroatoms; and 3) each ring in the system contains 3 to 7 ring members. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl, heteroarylalkoxy and the like) group may contain one or more substituents. Suitable substituents on the unsaturated carbon atom of an aryl, heteroaryl, aralkyl, or heteroaralkyl group are selected from halogen; haloalkyl;  $-\text{CF}_3$ ;  $-\text{R}^0$ ;  $-\text{OR}^0$ ;  $-\text{SR}^0$ ; 1,2-methylenedioxy; 1,2-ethylenedioxy; protected OH (such as acyloxy);

- phenyl (Ph); Ph substituted with  $R^0$ ;  $-O(Ph)$ ;  $-O(Ph)$  substituted with  $R^0$ ;  $-CH_2(Ph)$ ;  $-CH_2(Ph)$  substituted with  $R^0$ ;  $-CH_2CH_2(Ph)$ ;  $-CH_2CH_2(Ph)$  substituted with  $R^0$ ;  $-NO_2$ ;  $-CN$ ;  $-N(R^0)_2$ ;  $-NR^0C(O)R^0$ ;  $-NR^0C(O)N(R^0)_2$ ;  $-NR^0CO_2R^0$ ;  $-NR^0NR^0C(O)R^0$ ;  
 5  $-NR^0NR^0C(O)N(R^0)_2$ ;  $-NR^0NR^0CO_2R^0$ ;  $-C(O)C(O)R^0$ ;  $-C(O)CH_2C(O)R^0$ ;  $-CO_2R^0$ ;  $-C(O)R^0$ ;  $-C(O)N(R^0)_2$ ;  $-OC(O)N(R^0)_2$ ;  $-S(O)_2R^0$ ;  $-SO_2N(R^0)_2$ ;  $-S(O)R^0$ ;  $-NR^0SO_2N(R^0)_2$ ;  $-NR^0SO_2R^0$ ;  $-C(=S)N(R^0)_2$ ;  $-C(=NH)-N(R^0)_2$ ;  $-(CH_2)_yNHC(O)R^0$ ;  $-(CH_2)_yR^0$ ;  $-(CH_2)_yNHC(O)NHR^0$ ;  $-(CH_2)_yNHC(O)OR^0$ ;  $-(CH_2)_yNHS(O)R^0$ ;  
 10  $-(CH_2)_yNHCO_2R^0$ ;  $-(CH_2)_yNHC(O)CH(V_z-R^0)(R^0)$ ; wherein each  $R^0$  is independently selected from H, optionally substituted  $C_{1-6}$  aliphatic, an unsubstituted 5-10 membered heteroaryl or heterocyclic ring, phenyl (Ph),  $-O(Ph)$ , or  $-CH_2(Ph)$ ; wherein y is 0-6; z is 0-1; and V is a linker group.  
 15 When  $R^0$  is  $C_{1-6}$  aliphatic, it may be substituted with one or more substituents selected from  $-NH_2$ ,  $-NH(C_{1-4}$  aliphatic),  $-N(C_{1-4}$  aliphatic) $_2$ ,  $-S(O)(C_{1-4}$  aliphatic),  $-SO_2(C_{1-4}$  aliphatic), halogen,  $-C_{1-4}$  aliphatic,  $-OH$ ,  $-O(C_{1-4}$  aliphatic), nitro, cyano,  $-CO_2H$ ,  
 20  $-CO_2(C_{1-4}$  aliphatic),  $-O(halo\ C_{1-4}\ aliphatic)$ , or  $-halo(C_{1-4}$  aliphatic); wherein each  $C_{1-4}$  aliphatic is unsubstituted.

An aliphatic group or a non-aromatic heterocyclic ring may contain one or more substituents. Suitable substituents on the saturated carbon of an  
 25 aliphatic group or of a non-aromatic heterocyclic ring are selected from those listed above for the unsaturated carbon of an aryl or heteroaryl group and the following:  
 $=O$ ,  $=S$ ,  $=NNHR^9$ ,  $=NN(R^9)_2$ ,  $=N-$ ,  $OR^9$ ,  $=NNHC(O)R^9$ ,  $=NNHCO_2(alkyl)$ ,  $=NNHSO_2(alkyl)$ , or  $=NR^9$ , where each  $R^9$  is  
 30 independently selected from hydrogen or an optionally substituted  $C_{1-6}$  aliphatic group. When  $R^9$  is  $C_{1-6}$  aliphatic, it may be substituted with one or more substituents selected from amino, halogen, nitro, cyano, carboxy, t-butoxy, methoxy, ethoxy, hydroxy, or  $CF_3$ .

Substituents on the nitrogen of a non-aromatic heterocyclic ring are selected from  $-R^{10}$ ,  $-N(R^{10})_2$ ,  $-C(O)R^{10}$ ,  $-CO_2R^{10}$ ,  $-C(O)C(O)R^{10}$ ,  $-C(O)CH_2C(O)R^{10}$ ,  $-SO_2R^{10}$ ,  $-SO_2N(R^{10})_2$ ,  $-C(=S)N(R^{10})_2$ ,  $-C(=NH)-N(R^{10})_2$ , and  $-NR^{10}SO_2R^{10}$ ;  
5 wherein each  $R^{10}$  is independently selected from H, an optionally substituted  $C_{1-6}$  aliphatic, optionally substituted phenyl (Ph), optionally substituted  $-O(Ph)$ , optionally substituted  $CH_2(Ph)$ , optionally substituted  $-CH_2CH_2(Ph)$ , or an unsubstituted 5-6 membered heteroaryl  
10 or heterocyclic ring. When  $R^{10}$  is a  $C_{1-6}$  aliphatic group or a phenyl ring, it may be substituted with one or more substituents selected from  $NH_2$ ,  $NH(C_{1-4} \text{ aliphatic})$ ,  $-N(C_{1-4} \text{ aliphatic})_2$ , halogen,  $-C_{1-4} \text{ aliphatic}$ ,  $-OH$ ,  $-O-(C_{1-4} \text{ aliphatic})$ , nitro, cyano,  $-CO_2H$ ,  
15  $-CO_2(C_{1-4} \text{ aliphatic})$ ,  $-O(\text{halo } C_{1-4} \text{ aliphatic})$ , or  $-\text{halo}(C_{1-4} \text{ aliphatic})$ , wherein each  $C_{1-4}$  aliphatic is unsubstituted.

The term "linker group" or "linker" means an organic moiety that connects two parts of a compound.  
20 Linkers are comprised of  $-O-$ ,  $-S-$ ,  $-NH-$ ,  $-CH_2-$ ,  $-CO-$ , or an alkylidene chain. The alkylidene chain is a saturated or unsaturated, straight or branched,  $C_{1-6}$  carbon chain which is optionally substituted, and wherein up to two non-adjacent saturated carbons of the chain are  
25 optionally replaced by  $-C(O)-$ ,  $-C(O)C(O)-$ ,  $-C(O)NR^*-$ ,  $-C(O)NR^*NR^*-$ ,  $-CO_2-$ ,  $-OC(O)-$ ,  $-NR^*CO_2-$ ,  $-O-$ ,  $-NR^*C(O)NR^*-$ ,  $-OC(O)NR^*-$ ,  $-NR^*NR^*-$ ,  $-NR^*C(O)-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-NR^*-$ ,  $-SO_2NR^*-$ , or  $-NR^*SO_2-$ ; wherein  $R^*$  is selected from hydrogen or  $C_{1-4}$  aliphatic. Optional substituents on the  
30 alkylidene chain are as described above for an aliphatic group.

As used herein, linker group Q, when present, connects Ht with  $R^4$ . Q may form additional interactions within the ERK binding site to further enhance the

inhibitory activity of the compound. When Q is a carbonyl-containing moiety such as -C(O)-, -CO<sub>2</sub>-, -OC(O)-, -C(O)C(O)-, -C(O)NH-, -CO<sub>2</sub>NH-, -C(O)NHNH-, -NHC(O)-, -OC(O)NH-, or -NHCO<sub>2</sub>-, or a sulfonyl-containing moiety such as -SO<sub>2</sub>-, -SO<sub>2</sub>NH-, or -NHSO<sub>2</sub>-, the carbonyl or sulfonyl oxygen forms a hydrogen-bond with lysine 54 in the ERK binding site. When Q is an NH-containing moiety such as -CH<sub>2</sub>NH- or -NHNH-, the NH-group forms a hydrogen-bond with aspartic acid residue 167 in the ERK binding site. When Q is a hydrophobic group such as an alkyl chain, -O-, or -S-, Q forms additional hydrophobic interactions within the ERK binding site.

R<sup>4</sup> forms hydrophobic interactions within the binding site of ERK, especially with the side-chain carbons of lysine 54 and aspartic acid 167. R<sup>4</sup> may also form hydrophobic interactions with the glycine-rich loop which is made up of amino-acid residues 33-38. When R<sup>4</sup> is substituted, the substituents may form further interactions within the binding site to enhance the inhibitory activity of the compound. For example, when a substituent on R<sup>4</sup> is a hydrogen-bond donor or a hydrogen-bond acceptor, said substituent forms a hydrogen bond with enzyme-bound water molecules that exist in the binding site.

As used herein, linker group T, when present, connects the triazole ring with R<sup>2</sup>. T may form additional interactions within the ERK binding site to further enhance the inhibitory activity of the compound. When T is a carbonyl-containing moiety such as -C(O)-, -CO<sub>2</sub>-, -OC(O)-, -C(O)C(O)-, -C(O)NH-, -CO<sub>2</sub>NH-, -C(O)NHNH-, -NHC(O)-, -OC(O)NH-, or -NHCO<sub>2</sub>-, or a sulfonyl-containing moiety such as -SO<sub>2</sub>-, -SO<sub>2</sub>NH-, or -NHSO<sub>2</sub>-, the carbonyl or sulfonyl oxygen forms a hydrogen-bond with the NH of glutamine 105 in the ERK binding site. When T is an NH-



containing moiety such as  $-\text{CH}_2\text{NH}-$  or  $-\text{NHNH}-$ , the NH-group forms a hydrogen-bond with the carbonyl of glutamine 105. When T is a hydrophobic group such as an alkyl chain, -O- or -S-, Q forms additional hydrophobic interactions with the side-chain carbons of glutamine 105 as well as isoleucine 84.


The binding interactions described herein between the compounds of this invention and the ERK binding site have been determined by molecular modeling programs that are known to those of ordinary skill in the art. These molecular modeling programs include QUANTA [Molecular Simulations, Inc., Burlington, Mass., 1992] and SYBYL [Molecular Modeling Software, Tripos Associates, Inc., St. Louis, Mo., 1992]. As used herein, the amino acid numbering for the ERK enzyme corresponds to the Swiss-Prot database entry for accession #P28482. The Swiss-Prot database is an international protein sequence database distributed by the European Bioinformatics Institute (EBI) in Geneva, Switzerland. The database can be found at [www.ebi.ac.uk/swissprot](http://www.ebi.ac.uk/swissprot).

It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms, all such tautomeric forms of the compounds being within the scope of the invention.

Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the


present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -enriched carbon are within the scope of this invention.

5 One embodiment of this invention relates to a compound of formula I:




or a pharmaceutically acceptable salt or derivative thereof, wherein Ht,  $\text{R}^2$ , T, and m, n,  $\text{R}^3$ , Q and  $\text{R}^4$  are as described above, provided that when Ht is  
 10 tetrazol-5-yl, and m is 0,  $\text{R}^2$  is other than H or an unsubstituted phenyl; when m is 0 and  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{Q}_n\text{-R}^4$  are all H, Ht is other than [1,2,3]triazole; and when  $\text{T}_m\text{-R}^2$  and  $\text{Q}_n\text{-R}^4$  are both  $\text{C}(\text{O})\text{OH}$ , Ht is other than [1,2,3]triazol-5-yl.


15 Another embodiment of this invention relates to compounds wherein the Ht ring is a pyrrol-3-yl (II-A), [1,2,4]triazol-3-yl (II-B), [1,2,3]triazol-4-yl (II-C and II-D), or tetrazol-5-yl (II-E) ring as shown below:




II-A




II-B



II-C



II-D



II-E

or a pharmaceutically acceptable salt or  
 5 derivative thereof, wherein  $R^2$ , T, m, n,  $R^3$ , Q and  $R^4$  are as described above, provided that when Ht is tetrazol-5-yl, and  $m = 0$ ,  $R^2$  is other than H or an unsubstituted phenyl; when  $m = 0$  and  $R^2$ ,  $R^3$  and  $Q-R^4$  are all H, Ht is other than [1,2,3]triazole; when  $T-R^2$  and  $Q-R^4$  are both  
 10  $C(O)OH$ , Ht is other than [1,2,3]triazol-5-yl.

Preferred  $T_m R^2$  groups on II-A, II-B, II-C, II-D, and II-E are selected from hydrogen,  $N(R^7)_2$ , OH, 3-6 membered carbocyclyl, or an optionally substituted group selected from  $C_{1-6}$  aliphatic or a 5-6 membered aryl or  
 15 heteroaryl ring. When  $R^2$  is an optionally substituted phenyl or aliphatic group, preferred substituents on the phenyl or aliphatic group are methyl, ethyl, halo, nitro, alkoxy, and amino. Preferred  $T_m R^2$  groups are methyl, ethyl, propyl, cyclopropyl, cyclohexyl,  $CH_2OCH_3$ ,  $CH_2OH$ ,  
 20  $NH_2$ ,  $NHCH_3$ ,  $NHAc$ ,  $NHC(O)NHCH_3$ , and  $CH_2NHCH_3$ . More preferred  $T_m R^2$  groups of formulae II-A, II-B, II-C, II-D, and II-E are those listed in the Tables below.

When  $R^4$  is  $R^6$ , preferred  $R^6$  groups are pyrrolidin-1-yl, morpholin-4-yl, piperidin-1-yl, and  
 25 piperazin-1-yl, 4-methyl[1,4]diazepan-1-yl, 4-phenylpiperazine-1-yl, wherein each group is optionally substituted as described above for heterocyclic rings. When  $R^4$  is  $(CH_2)_y R^6$ ,  $(CH_2)_y CH(R^6)_2$ , or  $-N(R^5)_2$ , preferred  $R^6$  groups are pyridin-3-yl, pyridin-4-yl, imidazolyl, furan-  
 30 2-yl, 1,2,3,4-tetrahydroisoquinoline, tetrahydrofuran-2-yl, cyclohexyl, phenyl, benzyl,  $-CH_2OH$ ,  $-(CH_2)_2OH$ ,

isopropyl,  $-\text{CH}_2\text{NH}_2$ , and  $-(\text{CH}_2)_2\text{NH}_2$ , wherein each group is optionally substituted. Preferred substituents on  $\text{R}^6$  are  $-\text{OH}$ , pyridyl, piperidiny, and optionally substituted phenyl, wherein phenyl is optionally substituted as described above for aryl rings. When  $\text{R}^4$  is  $-(\text{CH}_2)_y\text{CH}(\text{R}^8)\text{CH}(\text{R}^6)_2$ , preferred  $\text{R}^8$  groups are  $\text{R}^7$  and  $\text{OR}^7$  such as  $\text{OH}$  and  $\text{CH}_2\text{OH}$ . Other preferred  $-\text{QR}^4$  groups are those listed in the Tables below.

Preferred compounds of formulae II-A, II-B, II-C, II-D, and II-E are those having one or more, more preferably more than one, and most preferably all, of the features selected from the group consisting of:

- (a)  $\text{T}_m\text{R}^2$  is hydrogen,  $\text{N}(\text{R}^7)_2$ ,  $\text{OH}$ , 3-6 membered carbocyclyl, or an optionally substituted group selected from  $\text{C}_{1-6}$  aliphatic or a 5-6 membered aryl or heteroaryl ring;
- (b)  $\text{Q}$  is  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{CONH}-$ ,  $-\text{SO}_2-$ ,  $-\text{SO}_2\text{NH}-$ ,  $-\text{OC}(\text{O})\text{NH}-$ ,  $-\text{C}(\text{O})\text{ONH}-$ , or  $-\text{CONHNH}-$ ;
- (c)  $\text{R}^4$  is  $-\text{NR}^5(\text{CH}_2)_y\text{N}(\text{R}^5)_2$ ,  $-(\text{CH}_2)_y\text{R}^6$ ,  $-(\text{CH}_2)_y\text{CH}(\text{R}^6)_2$ , or  $-(\text{CH}_2)_y\text{CH}(\text{R}^8)\text{CH}(\text{R}^6)_2$ ;
- (d)  $\text{R}^5$  is  $\text{R}$ ,  $\text{R}^7$ , or  $-(\text{CH}_2)_y\text{CH}(\text{R}^6)_2$ ; and
- (e)  $\text{R}^6$  is an optionally substituted group selected from phenyl, 5-6 membered heteroaryl, or 5-6 membered heterocyclyl.


More preferred compounds of formulae II-A, II-B, II-C, II-D, and II-E are those having one or more, more preferably more than one, or most preferably all, of the features selected from the group consisting of:

- (a)  $\text{T}_m\text{R}^2$  is selected from optionally substituted phenyl, methyl, ethyl, propyl, cyclopropyl, cyclohexyl,  $\text{CH}_2\text{OCH}_3$ ,  $\text{CH}_2\text{OH}$ ,  $\text{OH}$ ,  $\text{NH}_2$ ,  $\text{NHCH}_3$ ,  $\text{NHAc}$ ,  $\text{NHC}(\text{O})\text{NHCH}_3$ , or  $\text{CH}_2\text{NHCH}_3$ ;
- (b)  $\text{Q}$  is  $-\text{CO}-$ ,  $-\text{CONH}-$ ,  $-\text{SO}_2-$ , or  $-\text{SO}_2\text{NH}-$ ;

(c)  $R^4$  is  $-(CH_2)_yR^6$ ,  $-(CH_2)_yCH(R^6)_2$ , or  $-(CH_2)_yCH(R^8)CH(R^6)_2$ , wherein  $R^8$  is OH or  $CH_2OH$ ; and

(d)  $R^6$  is  $-CH_2OH$ ,  $-(CH_2)_2OH$ , isopropyl,  $-CH_2NH_2$ ,  $-(CH_2)_2NH_2$ , or an optionally substituted group selected from pyrrolidin-1-yl, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, 4-methyl[1,4]diazepan-1-yl, 4-phenyl-piperazine-1-yl, pyridin-3-yl, pyridin-4-yl, imidazolyl, furan-2-yl, 1,2,3,4-tetrahydroisoquinoline, tetrahydrofuran-2-yl, cyclohexyl, phenyl, or benzyl.

Preferred compounds of formulae II-A, II-B, and II-D are those of formulae II-A', II-B', and II-D':



Preferred  $R^6$  groups of formulae II-A', II-B', and II-D' are optionally substituted 6-membered aryl, heteroaryl, and carbocyclic rings, such as phenyl, pyridyl, and cyclohexyl.

Preferred  $T_mR^2$  groups of formulae II-A', II-B', and II-D' are as described above for formulae II-A, II-B, and II-D.

Preferred compounds of formulae II-A', II-B', and II-D' are those having one, and more preferably both, of the features selected from the group consisting of:

(a)  $T_mR^2$  is hydrogen,  $N(R^7)_2$ , OH, 3-6 membered carbocyclyl, or an optionally substituted group


selected from C<sub>1-6</sub> aliphatic or a 5-6 membered aryl or heteroaryl ring; and

- (b) R<sup>6</sup> is an optionally substituted 6-membered aryl, heteroaryl, or carbocyclic ring.

5 More preferred compounds of formulae II-A', II-B', and II-D' are those having one, and more preferably both, of the features selected from the group consisting of:

- 10 (a) T<sub>m</sub>R<sup>2</sup> is selected from optionally substituted phenyl, methyl, ethyl, propyl, cyclopropyl, cyclohexyl, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>OH, OH, NH<sub>2</sub>, NHCH<sub>3</sub>, NHAc, NHC(O)NHCH<sub>3</sub>, or CH<sub>2</sub>NHCH<sub>3</sub>; and
- (b) R<sup>6</sup> is cyclohexyl or an optionally substituted phenyl or pyridyl ring.

Preferred compounds of formulae II-A, II-B, and II-D are further selected from those of formulae II-A°, II-B°, and II-D°:



- 15 Preferred R<sup>6</sup> groups of formulae II-A°, II-B°, and II-D° are R or OR<sup>7</sup>. Examples of such groups include OH, CH<sub>2</sub>OH, or optionally substituted 6-membered aryl, heteroaryl, and carbocyclic rings, such as phenyl, pyridyl, and cyclohexyl. Preferred R<sup>8</sup> groups of formulae
- 20 II-A°, II-B°, and II-D° are R and OR<sup>7</sup>, wherein R is an optionally substituted group selected from C<sub>1-4</sub> aliphatic, 3-6 membered heterocyclic, or a 5-6 membered aryl or

heteroaryl ring. Examples of such groups include phenyl, methyl, ethyl, OH, and CH<sub>2</sub>OH.

Preferred T<sub>m</sub>R<sup>2</sup> groups of formulae II-A°, II-B°, and II-D° are as described above for formulae II-A, II-B,  
5 and II-D.

Preferred compounds of formulae II-A°, II-B°, and II-D° are those having one, and more preferably both, of the features selected from the group consisting of:

- (a) T<sub>m</sub>R<sup>2</sup> is hydrogen, N(R<sup>7</sup>)<sub>2</sub>, OH, 3-6 membered  
10 carbocyclyl, or an optionally substituted group selected from C<sub>1-6</sub> aliphatic or a 5-6 membered aryl or heteroaryl ring; and  
(b) R<sup>6</sup> is R or OR<sup>7</sup>, and R<sup>8</sup> is R<sup>7</sup> or OR<sup>7</sup>.

More preferred compounds of formulae II-A°, II-B°, and II-D° are those having one, and more preferably  
15 both, of the features selected from the group consisting of:

- (a) T<sub>m</sub>R<sup>2</sup> is selected from optionally substituted phenyl, methyl, ethyl, propyl, cyclopropyl, cyclohexyl, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>OH, OH, NH<sub>2</sub>, NHCH<sub>3</sub>, NHAc,  
20 NHC(O)NHCH<sub>3</sub>, or CH<sub>2</sub>NHCH<sub>3</sub>; and  
(b) R<sup>6</sup> is OH, CH<sub>2</sub>OH, phenyl, pyridyl, or cyclohexyl, and R<sup>8</sup> is methyl, ethyl, OH, or CH<sub>2</sub>OH.



Exemplary structures of formulae II-A, II-B, and II-D, wherein R<sup>3</sup> is H, m is zero, and n is one are set forth in Table 1 below.





25 Table 1. Compounds II-A, II-B, and II-D





No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-1, II-B-1, and II-C-1	phenyl	CON(Me) <sub>2</sub>
II-A-2, II-B-2, and II-C-2	phenyl	CO <sub>2</sub> Et
II-A-3, II-B-3, and II-C-3	3-NO <sub>2</sub> -phenyl	CONHNH <sub>2</sub>

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-4, II-B-4, and II-C-4	phenyl	CO (pyrrolidin-1-yl)
II-A-5, II-B-5, and II-C-5	phenyl	CONHCH <sub>2</sub> (Ph)
II-A-6, II-B-6, and II-C-6	3-NO <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-7, II-B-7, and II-C-7	4-Cl-phenyl	CO <sub>2</sub> Et
II-A-8, II-B-8, and II-C-8	4-OMe-phenyl	CO <sub>2</sub> Et
II-A-9, II-B-9, and II-C-9	3-NH <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-10, II-B-10, and II-C-10	3-OMe-phenyl	CO <sub>2</sub> Et
II-A-11, II-B-11, and II-C-11	4-F-phenyl	CO <sub>2</sub> Et
II-A-12, II-B-12, and II-C-12	4-NO <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-13, II-B-13, and II-C-13	3-Cl-phenyl	CO <sub>2</sub> Et
II-A-14, II-B-14, and II-C-14	3-F-phenyl	CO <sub>2</sub> Et
II-A-15, II-B-15, and II-C-15	phenyl	CO <sub>2</sub> H
II-A-16, II-B-16, and II-C-16	4-NH <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-17, II-B-17, and II-C-17	phenyl	CONHCH <sub>2</sub> CH <sub>2</sub> N (Me) <sub>2</sub>
II-A-18, II-B-18, and II-C-18	phenyl	CONHCH <sub>2</sub> (pyridin-3-yl)
II-A-19, II-B-19, and II-C-19	phenyl	CO (morpholin-4-yl)
II-A-20, II-B-20, and II-C-20	phenyl	CONH (isopropyl)
II-A-21, II-B-21, and II-C-21	phenyl	CO (4-Me-piperazin-1-yl)
II-A-22, II-B-22, and II-C-22	phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-23, II-B-23, and II-C-23	3-OMe-phenyl	CONMe <sub>2</sub>
II-A-24, II-B-24, and II-C-24	3-OMe-phenyl	CO (pyrrolidin-1-yl)
II-A-25, II-B-25, and II-C-25	3-OMe-phenyl	CONHCH <sub>2</sub> CH <sub>2</sub> N (Me) <sub>2</sub>
II-A-26, II-B-26, and II-C-26	3-OMe-phenyl	CONHCH <sub>2</sub> (pyridin-3-yl)
II-A-27, II-B-27, and II-C-27	3-OMe-phenyl	CO (morpholin-4-yl)
II-A-28, II-B-28, and II-C-28	3-OMe-phenyl	CONH (isopropyl)
II-A-29, II-B-29, and II-C-29	3-OMe-phenyl	CO (4-Me-piperazin-1-yl)
II-A-30, II-B-30, and II-C-30	3-OMe-phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-31, II-B-31, and II-C-31	4-NH <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-32, II-B-32, and II-C-32	H	CONMe <sub>2</sub>
II-A-33, II-B-33, and II-C-33	H	CO (pyrrolidin-1-yl)












No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-34, II-B-34, and II-C-34	3-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-35, II-B-35, and II-C-35	4-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-36, II-B-36, and II-C-36	3-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-37, II-B-37, and II-C-37	4-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-38, II-B-38, and II-C-38	3-Cl-phenyl	CONHBn
II-A-39, II-B-39, and II-C-39	3,5-dichlorophenyl	
II-A-40, II-B-40, and II-C-40	3-Br-phenyl	CONH(3,4-difluorophenyl)
II-A-41, II-B-41, and II-C-41	3-Cl-phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-42, II-B-42, and II-C-42	4-OH-3-I-5-nitrophenyl	CONH(2-OH-1-Ph-ethyl)
II-A-43, II-B-43, and II-C-43	3-Br-phenyl	
II-A-44, II-B-44, and II-C-44	5-NH <sub>2</sub> -4-OH-3-I-phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-45, II-B-45, and II-C-45	3-Br-phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-46, II-B-46, and II-C-46	3-Br-phenyl	CONHCH <sub>2</sub> (3-MeO-phenyl)
II-A-47, II-B-47, and II-C-47	3-Br-phenyl	CONHCH <sub>2</sub> (3-CF <sub>3</sub> -phenyl)
II-A-48, II-B-48, and II-C-48	3,5-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-49, II-B-49, and II-C-49	3-CF <sub>3</sub> -phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-50, II-B-50, and II-C-50	3-Cl-phenyl	CONHCH <sub>2</sub> Ph
II-A-51, II-B-51, and II-C-51	3,5-dichlorophenyl	CONHOCH <sub>2</sub> Ph
II-A-52, II-B-52, and II-C-52	4-OH-3-I-5-nitrophenyl	CONHCH <sub>2</sub> Ph
II-A-53, II-B-53, and II-C-53	3-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-54, II-B-54, and II-C-54	3,4-dichlorophenyl	CONHOCH <sub>2</sub> Ph
II-A-55, II-B-55, and II-C-55	3-Br-phenyl	CONHCH <sub>2</sub> (4-SO <sub>2</sub> Me-phenyl)
II-A-56, II-B-56, and II-C-56	3-Br-phenyl	CONHNH(3-CF <sub>3</sub> -phenyl)
II-A-57, II-B-57, and II-C-57	3-Cl-phenyl	CONHOCH <sub>2</sub> Ph



No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-58, II-B-58, and II-C-58	3-Br-phenyl	
II-A-59, II-B-59, and II-C-59	3-Br-phenyl	
II-A-60, II-B-60, and II-C-60	3-Br-phenyl	CONHCH <sub>2</sub> (2-Me-phenyl)
II-A-61, II-B-61, and II-C-61	3,4-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-62, II-B-62, and II-C-62	3-Br-phenyl	CONH (1-Ph-propyl)
II-A-63, II-B-63, and II-C-63	3-F-phenyl	CONHCH <sub>2</sub> Ph
II-A-64, II-B-64, and II-C-64	3,4-dichlorophenyl	
II-A-65, II-B-65, and II-C-65	3-Br-phenyl	
II-A-66, II-B-66, and II-C-66	3,5-dichlorophenyl	CON (Me) (Et)
II-A-67, II-B-67, and II-C-67	3-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-68, II-B-68, and II-C-68	3-Br-phenyl	CONHCH <sub>2</sub> (3,5-dimethoxyphenyl)
II-A-69, II-B-69, and II-C-69	3-Br-phenyl	CONHCH <sub>2</sub> (2-OMe-phenyl)
II-A-70, II-B-70, and II-C-70	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-71, II-B-71, and II-C-71	3-Cl-4-F-phenyl	CON (Me) (Et)
II-A-72, II-B-72, and II-C-72	3-Br-phenyl	CONH (2-OH-1-Ph-ethyl)
II-A-73, II-B-73, and II-C-73	3-NH <sub>2</sub> -phenyl	CONHCH <sub>2</sub> Ph
II-A-74, II-B-74, and II-C-74	3,4-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-75, II-B-75, and II-C-75	3-Me-phenyl	CONH (2-OH-1-Ph-ethyl)
II-A-76, II-B-76, and II-C-76	3,5-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-77, II-B-77, and II-C-77	3-Cl-4-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-78, II-B-78, and II-C-78	3,5-dichlorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-79, II-B-79, and II-C-79	3-NO <sub>2</sub> -phenyl	CONHCH <sub>2</sub> Ph

No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-80, II-B-80, and II-C-80	3-F-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-81, II-B-81, and II-C-81	3-Cl-2-F-phenyl	CON(Me)(Et)
II-A-82, II-B-82, and II-C-82	3-Cl-2-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-83, II-B-83, and II-C-83	3-Br-phenyl	
II-A-84, II-B-84, and II-C-84	3-Cl-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-85, II-B-85, and II-C-85	3,4-difluorophenyl	CONHOCH <sub>2</sub> Ph
II-A-86, II-B-86, and II-C-86	3-Br-phenyl	CONH(3-OH-1-Ph-propyl)
II-A-87, II-B-87, and II-C-87	3-Br-phenyl	
II-A-88, II-B-88, and II-C-88	3,4-difluorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-89, II-B-89, and II-C-89	3-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-90, II-B-90, and II-C-90	3-Me-phenyl	CONHCH <sub>2</sub> Ph
II-A-91, II-B-91, and II-C-91	3-Br-phenyl	
II-A-92, II-B-92, and II-C-92	4-Cl-phenyl	CONHCH <sub>2</sub> Ph
II-A-93, II-B-93, and II-C-93	3-Cl-phenyl	CON(Me)(Et)
II-A-94, II-B-94, and II-C-94	3-Br-phenyl	CONHCH <sub>2</sub> (4-SO <sub>2</sub> NH <sub>2</sub> -phenyl)
II-A-95, II-B-95, and II-C-95	3-OH-phenyl	CONHCH <sub>2</sub> Ph
II-A-96, II-B-96, and II-C-96	3-Me-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-97, II-B-97, and II-C-97	Phenyl	CONHCH <sub>2</sub> Ph
II-A-99, II-B-98, and II-C-98	2,5-difluorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-99, II-B-99, and II-C-99	4-Cl-phenyl	CONHOCH <sub>2</sub> Ph
II-A-100, II-B-100, and II-C-100	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-101, II-B-101, and II-C-101	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-102, II-B-102, and II-C-102	3-Br-phenyl	 CO(4-OH-4-Ph-piperidin-1-yl)

No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-103, II-B-103, and II-C-103	2,3-difluorophenyl	CONHOCH <sub>2</sub> Ph
II-A-104, II-B-104, and II-C-104	3-Cl-phenyl	CO (morpholin-4-yl)
II-A-105, II-B-105, and II-C-105	3-Br-phenyl	
II-A-106, II-B-106, and II-C-106	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-107, II-B-107, and II-C-107	3-Cl-4-F-phenyl	CO (morpholin-4-yl)
II-A-108, II-B-108, and II-C-108	3-Cl-4-F-phenyl	CON (Me) (Et)
II-A-109, II-B-109, and II-C-109	3-Br-phenyl	CONHCH <sub>2</sub> (4-NH <sub>2</sub> -phenyl)
II-A-110, II-B-110, and II-C-110	3-Br-phenyl	
II-A-111, II-B-111, and II-C-111	4-F-phenyl	CONHCH <sub>2</sub> Ph
II-A-112, II-B-112, and II-C-112	3,5-dichlorophenyl	CO (morpholin-4-yl)
II-A-113, II-B-113, and II-C-113	2,5-difluorophenyl	CONHOCH <sub>2</sub> Ph
II-A-114, II-B-114, and II-C-114	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-115, II-B-115, and II-C-115	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-116, II-B-116, and II-C-116	3,4-difluorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-117, II-B-117, and II-C-117	4-OMe-phenyl	CONHCH <sub>2</sub> Ph
II-A-118, II-B-118, and II-C-118	3-Br-phenyl	CONHCH <sub>2</sub> (2,4,6-trimethoxyphenyl)
II-A-119, II-B-119, and II-C-119	3-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-120, II-B-120, and II-C-120	3,4-difluorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-121, II-B-121, and II-C-121	3-Cl-2-F-phenyl	
II-A-122, II-B-122, and II-C-122	3-Br-phenyl	


No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-123, II-B-123, and II-C-123	3-Br-phenyl	
II-A-124, II-B-124, and II-C-124	3-Br-phenyl	CONHCH <sub>2</sub> (2,5-dimethoxyphenyl)
II-A-125, II-B-125, and II-C-125	3,5-dichlorophenyl	
II-A-126, II-B-126, and II-C-126	3-Br-phenyl	
II-A-127, II-B-127, and II-C-127	3,4-dichlorophenyl	CO(morpholin-4-yl)
II-A-128, II-B-128, and II-C-128	3-Br-phenyl	
II-A-129, II-B-129, and II-C-129	3-Cl-2-F-phenyl	CO(morpholin-4-yl)
II-A-130, II-B-130, and II-C-130	3-Br-phenyl	CONHCH <sub>2</sub> CH <sub>2</sub> OH
II-A-131, II-B-131, and II-C-131	3-NH <sub>2</sub> -phenyl	CONHCH <sub>2</sub> Ph
II-A-132, II-B-132, and II-C-132	3-MeOC(O)-phenyl	CONHCH <sub>2</sub> Ph
II-A-133, II-B-133, and II-C-133	4-MeO-phenyl	CONHOCH <sub>2</sub> Ph
II-A-134, II-B-134, and II-C-134	Phenyl	CO(pyrrolidin-1-yl)
II-A-135, II-B-135, and II-C-135	3-MeO-phenyl	CO(morpholin-4-yl)
II-A-136, II-B-136, and II-C-136	3-Cl-phenyl	CO(4-Me-piperidin-1-yl)
II-A-137, II-B-137, and II-C-137	3-NO <sub>2</sub> -phenyl	CONH <sub>2</sub> NH <sub>2</sub>
II-A-138, II-B-138, and II-C-138	3-Br-phenyl	
II-A-139, II-B-139, and II-C-139	3-Br-phenyl	
II-A-140, II-B-140, and II-C-140	3-Cl-phenyl	CONHPh
II-A-141, II-B-141, and II-C-141	2,3-difluorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-142, II-B-142, and II-C-142	3-Cl-phenyl	
II-A-143, II-B-143, and II-C-143	Phenyl	CON(Me) <sub>2</sub>

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-144, II-B-144, and II-C-144	3-OMe-phenyl	CO(pyrrolidin-1-yl)
II-A-145, II-B-145, and II-C-145	3-OMe-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-146, II-B-146, and II-C-146	4-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-147, II-B-147, and II-C-147	3-OMe-phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-148, II-B-148, and II-C-148	3-NO <sub>2</sub> -phenyl	COOEt
II-A-149, II-B-149, and II-C-149	Phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-150, II-B-150, and II-C-150	Phenyl	CO(morpholin-4-yl)
II-A-151, II-B-151, and II-C-151	3-Cl-phenyl	COOEt
II-A-152, II-B-152, and II-C-152	3-Br-phenyl	CONHMe
II-A-153, II-B-153, and II-C-153	Phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-154, II-B-154, and II-C-154	3-OMe-phenyl	CON(Me) <sub>2</sub>
II-A-155, II-B-155, and II-C-155	3-Cl-phenyl	
II-A-156, II-B-156, and II-C-156	3-Br-phenyl	
II-A-157, II-B-157, and II-C-157	3-Br-phenyl	COOEt
II-A-158, II-B-158, and II-C-158	phenyl	CONH(iPr)
II-A-159, II-B-159, and II-C-159	3-OMe-phenyl	CONH(iPr)
II-A-160, II-B-160, and II-C-160	3-COOH-phenyl	CONH(iPr)
II-A-161, II-B-161, and II-C-161	3-Br-phenyl	CONHO(iPr)
II-A-162, II-B-162, and II-C-162	3-F-phenyl	COOEt
II-A-163, II-B-163, and II-C-163	3-OMe-phenyl	CO(4-Me-piperidin-1-yl)
II-A-164, II-B-164, and II-C-164	4-NH <sub>2</sub> -phenyl	COOEt
II-A-165, II-B-165, and II-C-165	4-NO <sub>2</sub> -phenyl	COOEt
II-A-166, II-B-166, and II-C-166	phenyl	CO(4-Me-piperidin-1-yl)
II-A-167, II-B-167, and II-C-167	4-Cl-phenyl	COOEt
II-A-168, II-B-168, and II-C-168	4-OMe-phenyl	COOEt
II-A-169, II-B-169, and II-C-169	phenyl	COOEt
II-A-170, II-B-170, and II-C-170	3-OMe-phenyl	COOEt
II-A-171, II-B-171, and II-C-171	4-F-phenyl	COOEt

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-172, II-B-172, and II-C-172	3-NH <sub>2</sub> -phenyl	COOEt
II-A-173, II-B-173, and II-C-173	3-Cl-phenyl	COOH
II-A-174, II-B-174, and II-C-174	3-Cl-phenyl	
II-A-175, II-B-175, and II-C-175	3-Cl-phenyl	
II-A-176, II-B-176, and II-C-176	3-OMe-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-177, II-B-177, and II-C-177	3,5-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-178, II-B-178, and II-C-178	4-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-179, II-B-179, and II-C-179	4-OMe-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-180, II-B-180, and II-C-180	2,5-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-181, II-B-181, and II-C-181	2,5-difluorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-182, II-B-182, and II-C-182	4-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-183, II-B-183, and II-C-183	4-OMe-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-184, II-B-184, and II-C-184	3-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-185, II-B-185, and II-C-185	3-OMe-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-186, II-B-186, and II-C-186	2,5-dimethoxyphenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-187, II-B-187, and II-C-187	2,3-difluorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-188, II-B-188, and II-C-188	2,5-difluorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-189, II-B-189, and II-C-189	4-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-190, II-B-190, and II-C-190	4-OMe-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-191, II-B-191, and II-C-191	3-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-192, II-B-192, and II-C-192	3-OMe-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-193, II-B-193, and II-C-193	2,5-dimethoxyphenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-194, II-B-194, and II-C-194	3,4-difluorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-195, II-B-195, and II-C-195	2,3-difluorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-196, II-B-196, and II-C-196	2,5-difluorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-197, II-B-197, and II-C-197	4-F-phenyl	CO(morpholin-4-yl)
II-A-198, II-B-198, and II-C-198	4-OMe-phenyl	CO(morpholin-4-yl)
II-A-199, II-B-199, and II-C-199	3-F-phenyl	CO(morpholin-4-yl)
II-A-200, II-B-200, and II-C-200	2,5-dimethoxyphenyl	CO(morpholin-4-yl)
II-A-201, II-B-201, and II-C-201	3,4-difluorophenyl	CO(morpholin-4-yl)
II-A-202, II-B-202, and II-C-202	2,3-difluorophenyl	CO(morpholin-4-yl)
II-A-203, II-B-203, and II-C-203	2,5-difluorophenyl	CO(morpholin-4-yl)
II-A-204, II-B-204, and II-C-204	4-F-phenyl	CO(4-Me-piperidin-1-yl)
II-A-205, II-B-205, and II-C-205	4-OMe-phenyl	CO(4-Me-piperidin-1-yl)
II-A-206, II-B-206, and II-C-206	3-F-phenyl	CO(4-Me-piperidin-1-yl)
II-A-207, II-B-207, and II-C-207	2,5-dimethoxyphenyl	CO(4-Me-piperidin-1-yl)
II-A-208, II-B-208, and II-C-208	3,4-difluorophenyl	CO(4-Me-piperidin-1-yl)
II-A-209, II-B-209, and II-C-209	2,3-difluorophenyl	CO(4-Me-piperidin-1-yl)
II-A-210, II-B-210, and II-C-210	2,5-difluorophenyl	CO(4-Me-piperidin-1-yl)
II-A-211, II-B-211, and II-C-211	4-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-212, II-B-212, and II-C-212	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-213, II-B-213, and II-C-213	benzo[1,3]dioxo-5-yl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-214, II-B-214, and II-C-214	4-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-215, II-B-215, and II-C-215	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-216, II-B-216, and II-C-216	benzo[1,3]dioxo-5-yl	CONHCH <sub>2</sub> (pyrid-3-yl)










No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-217, II-B-217, and II-C-217	4-Cl-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-218, II-B-218, and II-C-218	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-219, II-B-219, and II-C-219	benzo [1,3] dioxo-5-yl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-220, II-B-220, and II-C-220	4-Cl-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-221, II-B-221, and II-C-221	3,4-dichlorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-222, II-B-222, and II-C-222	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-223, II-B-223, and II-C-223	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-224, II-B-224, and II-C-224	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-225, II-B-225, and II-C-225	benzo [1,3] dioxo-5-yl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-226, II-B-226, and II-C-226	3,5-dichlorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-227, II-B-227, and II-C-227	4-Cl-phenyl	CO (morpholin-4-yl)
II-A-228, II-B-228, and II-C-228	3,4-dimethoxyphenyl	CO (morpholin-4-yl)
II-A-229, II-B-229, and II-C-229	benzo [1,3] dioxo-5-yl	CO (morpholin-4-yl)
II-A-230, II-B-230, and II-C-230	4-Cl-phenyl	CO (4-Me-piperidin-1-yl)
II-A-231, II-B-231, and II-C-231	3,4-dichlorophenyl	CO (4-Me-piperidin-1-yl)
II-A-232, II-B-232, and II-C-232	3-Cl-2-F-phenyl	CO (4-Me-piperidin-1-yl)
II-A-233, II-B-233, and II-C-233	3-Cl-4-F-phenyl	CO (4-Me-piperidin-1-yl)
II-A-234, II-B-234, and II-C-234	3,4-dimethoxyphenyl	CO (4-Me-piperidin-1-yl)
II-A-235, II-B-235, and II-C-235	benzo [1,3] dioxo-5-yl	CO (4-Me-piperidin-1-yl)
II-A-236, II-B-236, and II-C-236	3,5-dichlorophenyl	CO (4-Me-piperidin-1-yl)
II-A-237, II-B-237, and II-C-237	2,3-difluorophenyl	CON (Me) (Et)
II-A-238, II-B-238, and II-C-238	4-F-phenyl	

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-239, II-B-239, and II-C-239	3-OMe-phenyl	
II-A-240, II-B-240, and II-C-240	2,5-dimethoxyphenyl	
II-A-241, II-B-241, and II-C-241	3,4-difluorophenyl	
II-A-242, II-B-242, and II-C-242	2,3-difluorophenyl	
II-A-243, II-B-243, and II-C-243	2,5-difluorophenyl	
II-A-244, II-B-244, and II-C-244	3-MeO-phenyl	CONHOCH <sub>2</sub> Ph
II-A-245, II-B-245, and II-C-245	2,5-dimethoxyphenyl	CONHOCH <sub>2</sub> Ph
II-A-246, II-B-246, and II-C-246	3-F-phenyl	
II-A-247, II-B-247, and II-C-247	3-MeO-phenyl	
II-A-248, II-B-248, and II-C-248	3,4-difluorophenyl	
II-A-249, II-B-249, and II-C-249	2,3-difluorophenyl	
II-A-250, II-B-250, and II-C-250	3-Cl-phenyl	
II-A-251, II-B-251, and II-C-251	4-Cl-phenyl	
II-A-252, II-B-252, and II-C-252	4-Cl-phenyl	
II-A-253, II-B-253, and II-C-253	3,4-dichlorophenyl	













No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-254, II-B-254, and II-C-254	3,4-dichlorophenyl	
II-A-255, II-B-255, and II-C-255	3-Cl-2-F-phenyl	
II-A-256, II-B-256, and II-C-256	3-Cl-4-F-phenyl	
II-A-257, II-B-257, and II-C-257	3-Cl-4-F-phenyl	
II-A-258, II-B-258, and II-C-258	3,4-dimethoxyphenyl	CON (Me) (Et)
II-A-259, II-B-259, and II-C-259	3,4-dimethoxyphenyl	
II-A-260, II-B-260, and II-C-260	3,4-dimethoxyphenyl	CONHOCH <sub>2</sub> Ph
II-A-261, II-B-261, and II-C-261	3,4-dimethoxyphenyl	
II-A-262, II-B-262, and II-C-262	benzo[1,3]dioxo-5-yl	CON (Me) (Et)
II-A-263, II-B-263, and II-C-263	benzo[1,3]dioxo-5-yl	
II-A-264, II-B-264, and II-C-264	benzo[1,3]dioxo-5-yl	CONHOCH <sub>2</sub> Ph
II-A-265, II-B-265, and II-C-265	benzo[1,3]dioxo-5-yl	
II-A-266, II-B-266, and II-C-266	3,5-dichlorophenyl	
II-A-267, II-B-267, and II-C-267	3-Br-phenyl	
II-A-268, II-B-268, and II-C-268	3-Br-phenyl	
II-A-269, II-B-269, and II-C-269	3-Br-phenyl	

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-270, II-B-270, and II-C-270	3-Br-phenyl	
II-A-271, II-B-271, and II-C-271	3-Br-phenyl	
II-A-272, II-B-272, and II-C-272	3-Br-phenyl	
II-A-273, II-B-273, and II-C-273	3-Br-phenyl	
II-A-274, II-B-274, and II-C-274	3-Br-phenyl	
II-A-275, II-B-275, and II-C-275	3-Br-phenyl	
II-A-276, II-B-276, and II-C-276	3-Br-phenyl	
II-A-277, II-B-277, and II-C-277	3-Br-phenyl	
II-A-278, II-B-278, and II-C-278	3-Br-phenyl	
II-A-279, II-B-279, and II-C-279	3-Br-phenyl	
II-A-280, II-B-280, and II-C-280	3-Br-phenyl	CONH(CH <sub>2</sub> ) <sub>2</sub> COOH
II-A-281, II-B-281, and II-C-281	3-Br-phenyl	
II-A-282, II-B-282, and II-C-282	3-Br-phenyl	CONHCH <sub>2</sub> (4-COOH-phenyl)
II-A-283, II-B-283, and II-C-283	3-Br-phenyl	





No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-284, II-B-284, and II-C-284	3-Br-phenyl	
II-A-285, II-B-285, and II-C-285	3-NO <sub>2</sub> -phenyl	CONHCH <sub>2</sub> phenyl
II-A-286, II-B-286, and II-C-286	3-Cl-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-287, II-B-287, and II-C-287	3-(N-Et-NHCO)-phenyl	CONHCH <sub>2</sub> phenyl
II-A-288, II-B-288, and II-C-288	3-Br-phenyl	
II-A-289, II-B-289, and II-C-289	3-NO <sub>2</sub> -phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-290, II-B-290, and II-C-290	3-Br-phenyl	
II-A-291, II-B-291, and II-C-291	3-F-phenyl	CON(Me)(Et)
II-A-292, II-B-292, and II-C-292	3-MeO-phenyl	CON(Me)(Et)
II-A-293, II-B-293, and II-C-293	3-Br-phenyl	
II-A-294, II-B-294, and II-C-294	3-Br-phenyl	
II-A-295, II-B-295, and II-C-295	3-Br-phenyl	
II-A-296, II-B-296, and II-C-296	3-Br-phenyl	
II-A-297, II-B-297, and II-C-297	phenyl	CONH(CH <sub>2</sub> ) <sub>2</sub> NMe <sub>2</sub>
II-A-298, II-B-298, and II-C-298	3-MeO-phenyl	CONH(CH <sub>2</sub> ) <sub>2</sub> NMe <sub>2</sub>
II-A-299, II-B-299, and II-C-299	3-Br-phenyl	CONHCH <sub>2</sub> phenyl

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-300, II-B-300, and II-C-300	3-Cl-phenyl	
II-A-301, II-B-301, and II-C-301	3-Cl-phenyl	
II-A-302, II-B-302, and II-C-302	3-Cl-phenyl	
II-A-303, II-B-303, and II-C-303	3-Cl-phenyl	
II-A-304, II-B-304, and II-C-304	3-Cl-phenyl	
II-A-305, II-B-305, and II-C-305	3-Cl-phenyl	
II-A-306, II-B-306, and II-C-306	3-Cl-phenyl	
II-A-307, II-B-307, and II-C-307	3-Cl-phenyl	
II-A-308, II-B-308, and II-C-308	3-Cl-phenyl	
II-A-309, II-B-309, and II-C-309	3-Cl-phenyl	
II-A-310, II-B-310, and II-C-310	3,5-dichlorophenyl	
II-A-311, II-B-311, and II-C-311	3-Br-5-CF <sub>3</sub> -phenyl	

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-312, II-B-312, and II-C-312	3-Cl-phenyl	
II-A-313, II-B-313, and II-C-313	3,5-dichlorophenyl	
II-A-314, II-B-314, and II-C-314	3-Cl-4-CN-phenyl	
II-A-315, II-B-315, and II-C-315	3-Cl-4-CH2OH-phenyl	
II-A-316, II-B-316, and II-C-316	3-Cl-4-CH2NH2-phenyl	
II-A-317, II-B-317, and II-C-317		
II-A-318, II-B-318, and II-C-318		
II-A-319, II-B-319, and II-C-319		
II-A-320, II-B-320, and II-C-320		
II-A-321, II-B-321, and II-C-321		
II-A-322, II-B-322, and II-C-322		
II-A-323, II-B-323, and II-C-323	CH <sub>2</sub> Ph	CON(Me) <sub>2</sub>








No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-324, II-B-324, and II-C-324	cyclopentylmeth yl	CO <sub>2</sub> NHCH <sub>2</sub> Ph
II-A-325, II-B-325, and II-C-325	isopropyl	CN
II-A-326, II-B-326, and II-C-326	3-Cl-phenyl	NHCOCH <sub>2</sub> Ph
II-A-327, II-B-327, and II-C-327	3-Cl-phenyl	NHSO <sub>2</sub> -morpholin-4- yl
II-A-328, II-B-328, and II-C-328	3-Cl-phenyl	NHCONHCH <sub>2</sub> Ph
II-A-329, II-B-329, and II-C-329	3-Cl-phenyl	NHCO <sub>2</sub> - tetrahydrofuran- 2-yl
II-A-330, II-B-330, and II-C-330	CH <sub>2</sub> Ph	CONHCH <sub>2</sub> Ph
II-A-331, II-B-331, and II-C-331	Me	CONHCH <sub>2</sub> Ph
II-A-332, II-B-332, and II-C-332	isopropyl	CONHCH <sub>2</sub> Ph
II-A-333, II-B-333, and II-C-333	H	CON (Me) <sub>2</sub>
II-A-334, II-B-334, and II-C-334		
II-A-335, II-B-335, and II-C-335		
II-A-336, II-B-336, and II-C-336		
II-A-337, II-B-337, and II-C-337		
II-A-338, II-B-338, and II-C-338		
II-A-339, II-B-339, and II-C-339		






























No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-340, II-B-340, and II-C-340		
II-A-341, II-B-341, and II-C-341		
















Additional preferred compounds, including those of formulae II-A', II-B', II-D', II-A°, II-B°, and II-D° are set forth in Table 2 below.

5 Table 2. Additional preferred compounds

No.	T <sub>m</sub> -R <sup>2</sup>	Q <sub>n</sub> -R <sup>4</sup>
II-A-342, II-B-342, and II-C-342	2-F-3-Cl-phenyl	
II-A-343, II-B-343, and II-C-343	methyl	
II-A-344, II-B-344, and II-C-344	methyl	
II-A-345, II-B-345, and II-C-345	Methyl	
II-A-346, II-B-346, and II-C-346	3,5-dichlorophenyl	
II-A-347, II-B-347, and II-C-347	3-F,5-CF <sub>3</sub> -phenyl	
II-A-348, II-B-348, and II-C-348	Methyl.	

No.	$T_m-R^2$	$Q_n-R^4$
II-A-349, II-B-349, and II-C-349	H	
II-A-350, II-B-350, and II-C-350	Methyl	
II-A-351, II-B-351, and II-C-351	Methyl	
II-A-352, II-B-352, and II-C-352	Methyl	
II-A-353, II-B-353, and II-C-353	Cyclohexyl	
II-A-354, II-B-354, and II-C-354	Cyclopropyl	
II-A-355, II-B-355, and II-C-355	Methyl	
II-A-356, II-B-356, and II-C-356	Methyl	
II-A-357, II-B-357, and II-C-357	$CH_2OCH_3$	
II-A-358, II-B-358, and II-C-358	$CH_2OH$	
II-A-359, II-B-359, and II-C-359	Methyl	
II-A-360, II-B-360, and II-C-360	Methyl	
II-A-361, II-B-361, and II-C-361	Methyl	
II-A-362, II-B-362, and II-C-362	Methyl	

No.	T <sub>m</sub> -R <sup>2</sup>	Q <sub>n</sub> -R <sup>4</sup>
II-A-363, II-B-363, and II-C-363	Methyl	
II-A-364, II-B-364, and II-C-364	H	
II-A-365, II-B-365, and II-C-365	H	
II-A-366, II-B-366, and II-C-366	H	
II-A-367, II-B-367, and II-C-367	H	
II-A-368, II-B-368, and II-C-368	H	
II-A-369, II-B-369, and II-C-369	H	
II-A-370, II-B-370, and II-C-370	Methyl	
II-A-371, II-B-371, and II-C-371	Methyl	
II-A-372, II-B-372, and II-C-372	Methyl	
II-A-373, II-B-373, and II-C-373	Methyl	
II-A-374, II-B-374, and II-C-374	Methyl	
II-A-375, II-B-375, and II-C-375	Methyl	







No.	T <sub>m</sub> -R <sup>2</sup>	Q <sub>n</sub> -R <sup>4</sup>
II-A-376, II-B-376, and II-C-376	Methyl	
II-A-377, II-B-377, and II-C-377	Methyl	
II-A-378, II-B-378, and II-C-378	Methyl	
II-A-379, II-B-379, and II-C-379	Methyl	
II-A-380, II-B-380, and II-C-380	methyl	
II-A-381, II-B-381, and II-C-381	H	
II-A-382, II-B-382, and II-C-382	H	
II-A-383, II-B-383, and II-C-383		
II-A-384, II-B-384, and II-C-384		
II-A-385, II-B-385, and II-C-385		
II-A-386, II-B-386, and II-C-386		







No.	$T_m-R^2$	$Q_n-R^4$
II-A-387, II-B-387, and II-C-387		
II-A-388, II-B-388, and II-C-388		
II-A-389, II-B-389, and II-C-389		
II-A-390, II-B-390, and II-C-390		







Additional exemplary compounds of formulae II-A, II-B, II-C, and II-E are set forth in Table 3 below.

5 Table 3. Compounds II-A, II-B, II-C, and II-E







No.	Structure
II-A 1	
II-A 2	
II-A 3	

No.	Structure
II-A 4	
II-A 5	
II-A 6	
II-A 7	
II-A 8	
II-B 1	

No.	Structure
II-B 2	 <chem>COC(=O)Oc1ccc(Cl)cc1C2=NN=C(C3=NC(=C(C4=NC(=C(C5=NC(=C(C6=CC(=CC(=C6)C(F)(F)F)CCNC(=O)N3)N4)N5)N6)N)N2)N1</chem>
II-B 3	 <chem>c1ccc(cc1)CC2=NN=C(C3=NC(=C(C4=NC(=C(C5=CC(=CC(=C5)Cl)CC2)N4)N3)N1)N5</chem>
II-B 4	 <chem>OCc1ccc(Cl)cc1C2=NN=C(C3=NC(=C(C4=NC(=C(C5=CC=CC=N5)CCNC(=O)N3)N4)N2)N1</chem>
II-B 5	 <chem>OC1=CC(=CC(=C1C(F)(F)F)C2=NN=C(C3=NC(=C(C4=NC(=C(C5=CC=CC=N5)CCNC(=O)N3)N4)N2)N1</chem>
II-B 6	 <chem>Fc1ccc(cc1F)CCNC2=NN=C(C3=NC(=C(C4=NC(=C(C5=CC(=CC(=N5)O)N4)N3)N2)N1</chem>
II-C 1	 <chem>CN(C)Cc1ccc(Cl)cc1C2=NN=C(C3=NC(=C(C4=NC(=C(C5=CC(=CC(=C5)C)CC2)N4)N3)N1</chem>







No.	Structure
II-C 2	
II-C 3	
II-C 4	
II-C 5	
II-C 6	
II-C 7	

















No.	Structure
II-E 1	
II-E 2	
II-E 3	
II-E 4	
II-E 5	
II-E 6	








Additional preferred compounds of formula II-A are provided in Table 4 below:




Table 4: Additional Compounds II-A

No.	Structure
II-A 9	 <chem>Clc1ccc(cc1)c2nn[nH]c2c3cc[nH]3CO</chem>
II-A 10	 <chem>Clc1ccc(cc1)c2nn[nH]c2c3cc[nH]3CN</chem>
II-A 11	 <chem>COc1cc[nH]c1c2nn[nH]2c3cc[nH]3C(=O)NCC(C)C</chem>
II-A 12	 <chem>c1ccc2c(c1)sc(n2)c3cc[nH]c3c4nn[nH]4c5cc[nH]5C(=O)NCC(C)C</chem>
II-A 13	 <chem>c1ccc2c(c1)c[nH]c2c3cc[nH]c3c4nn[nH]4c5cc[nH]5C(=O)NCC(C)C</chem>
II-A 14	 <chem>CC(C)C(=O)c1cc[nH]c1c2nn[nH]2c3cc[nH]3C(=O)NCC(C)C</chem>





No.	Structure
II-A 15	
II-A 16	
II-A 17	
II-A 18	
II-A 19	
II-A 20	
II-A 21	

No.	Structure
II-A 22	
II-A 23	
II-A 24	
II-A 25	
II-A 26	
II-A 27	
II-A 28	

No.	Structure
II-A 29	 <chem>O=C(c1ccccc1)c2cc[nH]c2c3nn[nH]c3c4cc[nH]4</chem>
II-A 30	 <chem>O=C(c1ccccc1)c2cc[nH]c2c3nn[nH]c3c4cc[nH]4CCN5CCOCC5</chem>
II-A 31	 <chem>NC(=O)c1cc(Cl)ccc1c2cc[nH]c2c3nn[nH]c3c4cc[nH]4CC5=CN=CC=C5</chem>
II-A 32	 <chem>O=C(c1cc[nH]c1)c2cc[nH]c2c3nn[nH]c3c4cc[nH]4CCOCC5=CC=CC=C5</chem>
II-A 33	 <chem>OCC1=CC=CC=C1c2cc[nH]c2c3nn[nH]c3c4cc[nH]4CCOCC5=CC=CC=C5</chem>
II-A 34	 <chem>O=C(NC1=CC=CC=C1)Oc2cc[nH]c2c3nn[nH]c3c4cc[nH]4CCOCC5=CC=CC=C5</chem>
II-A 35	 <chem>O=C(c1ccccc1)c2cc[nH]c2c3nn[nH]c3c4cc[nH]4CC5=CC=CC=C5</chem>

No.	Structure
II-A 36	
II-A 37	
II-A 38	

Preferred embodiments of this invention are represented by formulae **III-A**, **III-B**, **III-C**, and **III-D**:

**III-A****III-B****III-C****III-D,**

wherein T, m, R<sup>2</sup>, R<sup>4</sup> are as described above.

Preferred R<sup>2</sup> and R<sup>4</sup> of formulae **III-A**, **III-B**, **III-C**, and **III-D** are as described above for formulae **II-A**, **II-B**, **II-C**, and **II-D**.


5

Another object of the invention is to provide methods of producing the above-identified compounds of formula **I**. Schemes 1-6 depict syntheses of the triazole-pyrrole (Schemes 1-2), triazole-[1,2,4]triazole (Scheme 3), triazole-[1,2,3]triazole (Schemes 4-5), and triazole-

10

tetrazole (Scheme 6) compounds of this invention. One having ordinary skill in the art will recognize that Schemes 1-6 may be altered according to the functional groups present to make analogous compounds of this invention.


### Scheme 1 - Triazole-pyrrole



Reagents and conditions: (a) K<sub>2</sub>CO<sub>3</sub>, toluene, reflux

Scheme 1 above shows a general synthetic route used to prepare the pyrrol-3-yl compounds of formula II-A of this invention when m is zero and R<sup>2</sup> is an optionally substituted phenyl group. In Scheme 1, the triazole ring is formed by combining a cyano-1-pyrrole such as 1 and a diazo compound such as 2 in the presence of K<sub>2</sub>CO<sub>3</sub> in refluxing toluene. One of skill in the art will recognize that the above scheme may be used to synthesize compounds wherein R<sup>2</sup> is other than substituted phenyl. In addition, a compound such as 3 may be further derivatized to produce a variety of substituents Q<sub>n</sub>-R<sup>4</sup>.


### Scheme 2 - Triazole-pyrrole



Reagents and conditions: (a)  $\text{NaN}_3$ , DMSO

Scheme 2 shows an alternate method for synthesizing the pyrrol-3-yl compounds II-A of the present invention. In Scheme 2, the triazole ring is formed by reacting a compound such as pyrrole derivative 4 with  $\text{NaN}_3$  in DMSO. One of skill in the art will recognize that the above scheme may be used to synthesize compounds with a variety of  $\text{Q}_n\text{-R}^4$  substituents. In addition, a compound such as 5 may be further derivatized to produce a variety of substituents  $\text{T}_m\text{-R}^2$ .

Scheme 3 - Triazole-[1,2,4]triazole




Reagents and conditions: (a) acetic anhydride,  $140^\circ\text{C}$ ;  
(b).(i) 2,  $\text{KOtBu}$ , toluene, reflux, (ii)  $\text{H}_2$ ,  $\text{Pd/C}$ ,  $\text{EtOH}$

Scheme 3 depicts the [1,2,4]triazole formation in step (a) by the method described in Bladin et al., Chem. Ber., 1885, pp. 1544. Step (b) depicts formation of the [1,2,3]triazole ring as in Scheme 1 above. One of skill in the art will recognize that Scheme 3 may be used to synthesize compounds wherein  $\text{R}^2$  is other than substituted phenyl. In addition, a compound such as 7 may be further derivatized to produce a variety of substituents  $\text{Q}_n\text{-R}^4$ .

20


25



Scheme 4 - Triazole-[1,2,3]triazole


Reagents and conditions: (a) TMS-N<sub>3</sub>, NaN<sub>3</sub>, NaOH

5                    Scheme 4 depicts the cyclization to form the [1,2,3]triazole-[1,2,3]triazole system according to the method of Banert et al., Chem. Ber., 1989, pp. 1175. One of skill in the art will recognize that Scheme 4 may be used to synthesize compounds wherein R<sup>2</sup> is other than substituted phenyl. In addition, a compound such as 9  
 10                    may be further derivatized to produce a variety of substituents Q<sub>n</sub>-R<sup>4</sup>.

Scheme 5 - Triazole-[1,2,3]triazole

Reagents and conditions: (a) TMS-N<sub>3</sub>, NaN<sub>3</sub>, NaOH

                    Scheme 5 depicts another [1,2,3]triazole-[1,2,3]triazole cyclization according to the method of Banert et al. above. One of skill in the art will  
 20                    recognize that Scheme 5 may be used to synthesize compounds wherein R<sup>2</sup> is other than substituted phenyl. In addition, a compound such as 11 may be further derivatized to produce a variety of substituents Q<sub>n</sub>-R<sup>4</sup>.

Scheme 6 - Triazole-tetrazole

Reagents and conditions: (a) NaN<sub>3</sub>, DMF; (b) NaN<sub>3</sub>, NH<sub>4</sub>Cl, DMF

5

Scheme 6 depicts the formation of the triazole in step (a) according to the method of Beck et al., Chem. Ber., 1973, pp. 106 and the formation of the tetrazole in step (b) according to the method of Katritzky et al., J. Heterocycl. Chem., 1996, pp. 1107. One of skill in the art will recognize that the above scheme may be used to synthesize compounds wherein R<sup>2</sup> is other than substituted phenyl. In addition, a compound such as 14 may be further derivatized to produce a variety of substituents Q<sub>n</sub>-R<sup>4</sup>.

Schemes 1-6 above show general synthetic routes useful in preparing the compounds of this invention wherein T<sub>m</sub>-R<sup>2</sup> is an optionally substituted phenyl group, R<sup>3</sup> is H or methyl, and Q<sub>n</sub>-R<sup>4</sup> is methyl or a carbonyl derivative. One having skill in the art may synthesize other compounds of this invention following the teachings of this specification, using reagents that are readily synthesized or commercially available.

The activity of a compound utilized in this invention as an inhibitor of ERK, may be assayed *in vitro*, *in vivo* or in a cell line according to methods known in the art. *In vitro* assays include assays that determine inhibition of either the phosphorylation activity or ATPase activity of activated ERK. Alternate *in vitro* assays quantitate the ability of the inhibitor

to bind to ERK. Inhibitor binding may be measured by radiolabelling the inhibitor prior to binding, isolating the inhibitor/ERK complex and determining the amount of radiolabel bound. Alternatively, inhibitor binding may  
5 be determined by running a competition experiment where new inhibitors are incubated with ERK bound to known radioligands. Detailed conditions for assaying a compound utilized in this invention as an inhibitor of ERK kinase are set forth in the Examples below.

10           According to another embodiment, the invention provides a composition comprising a compound of this invention or a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier, adjuvant, or vehicle. The amount of compound in the  
15 compositions of this invention is such that is effective to detectably inhibit a protein kinase, particularly ERK in a biological sample or in a patient. Preferably the composition of this invention is formulated for administration to a patient in need of such composition.  
20 Most preferably, the composition of this invention is formulated for oral administration to a patient.

The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

The term "pharmaceutically acceptable carrier, adjuvant, or vehicle" refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the  
25 compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of  
30

saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

10           The term "detectably inhibit", as used herein means a measurable change in ERK activity between a sample comprising said composition and an ERK kinase and an equivalent sample comprising ERK kinase in the absence of said composition. According to a preferred  
15           embodiment, inhibition of kinase activity by a compound according to the present invention is greater than 10% compared to the kinase activity in the absence of the compound. Preferably, inhibition is greater than 20%, 30%, or 40%, and even more preferably greater than 50%,  
20           60%, 70%, 80%, or 90%.

          A "pharmaceutically acceptable derivative" means any non-toxic salt, ester, salt of an ester or other derivative of a compound of this invention that, upon administration to a recipient, is capable of  
25           providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.

          Pharmaceutically acceptable salts of the compounds of this invention include those derived from  
30           pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate,

digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., magnesium), ammonium and  $N^+(C_{1-4} \text{ alkyl})_4$  salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

Preferably, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art

using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added.

For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

Alternatively, the pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically

acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

For ophthalmic use, the pharmaceutically acceptable compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum.

The pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

Most preferably, the pharmaceutically acceptable compositions of this invention are formulated for oral administration.

The amount of the compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01 - 100



mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular compound in the composition.

Depending upon the particular condition, or disease, to be treated or prevented, additional therapeutic agents, which are normally administered to treat or prevent that condition, may also be present in the compositions of this invention. As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

For example, chemotherapeutic agents or other anti-proliferative agents may be combined with the compounds of this invention to treat proliferative diseases and cancer. Examples of known chemotherapeutic agents include, but are not limited to, Gleevec™, adriamycin, dexamethasone, vincristine, cyclophosphamide, fluorouracil, topotecan, taxol, interferons, and platinum derivatives.

Other examples of agents the compounds of this invention may also be combined with include, without limitation, anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine,

cyclophosphamide, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and  
5 sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers,  
10 ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and anti-viral agents; agents for treating blood disorders such as corticosteroids, anti-leukemic agents, and growth  
15 factors; agents for treating diabetes such as insulin, insulin analogues, alpha glucosidase inhibitors, biguanides, and insulin sensitizers; and agents for treating immunodeficiency disorders such as gamma globulin.

20           The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional  
25 therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

          According to another embodiment, the invention  
30 relates to a method of inhibiting ERK kinase activity in a biological sample comprising the step of contacting said biological sample with a compound of this invention, or a pharmaceutically acceptable composition comprising said compound.

The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

Inhibition of ERK kinase activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, blood transfusion, organ-transplantation, biological specimen storage, and biological assays.

According to another embodiment, the invention provides a method for treating or lessening the severity of an ERK-mediated disease or condition in a patient comprising the step of administering to said patient a pharmaceutically acceptable composition according to the present invention.

The term "ERK-mediated condition" or "disease", as used herein, means any disease or other deleterious condition in which ERK is known to play a role. The term "ERK-mediated condition" or "disease" also means those diseases or conditions that are alleviated by treatment with an ERK inhibitor. Such conditions include, without limitation, cancer, stroke, diabetes, hepatomegaly, cardiovascular disease including cardiomegaly, Alzheimer's disease, cystic fibrosis, viral disease, autoimmune diseases, atherosclerosis, restenosis, psoriasis, allergic disorders including asthma, inflammation, neurological disorders and hormone-related diseases. The term "cancer" includes, but is not limited to the following cancers: breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin, keratoacanthoma, lung, epidermoid carcinoma, large cell

carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma, bladder carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity and pharynx (oral), lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central nervous system, and leukemia.

Compounds of the present invention are also useful as inhibitors of related kinases to ERK. The term "related kinases" refer to protein kinases having residues which are similar to those residues which line the ERK binding site. Without wishing to be bound by theory, applicants speculate that this inhibitory activity is due to the close structural similarity between the active sites of ERK and related kinases. The alignment of the ERK sequence with other kinases can be derived from common software programs such as the "bestfit" program available from Genetics Computer Group. This program uses the local homology algorithm described by Smith and Waterman in *Advances in Applied Mathematics* 2; 482 (1981).

Related kinases inhibited by the compounds of this invention would contain residues, identified by the above standard protein sequence alignment software, corresponding to the ERK residues: I31, E33, G34, A35, Y36, G37, M38, V39, A52, K54, R67, T68, E71, L75, I84, I86, I103, Q105, D106, L107, M108, E109, D111, K114, D149, K151, S153, N154, L156, C166, and D167, with a similarity score of 80% or greater. In a more preferred embodiment the similarity score is 85%, more preferably 90%, even more preferably 95%, 96%, 97% or 98%. The

similarity score may be determined using standard amino acid substitution tables such as those described by Dayhoff (Dayhoff, M.O., et al, *Atlas of Protein Sequence and Structure*, 1979) and Blossom-Henikoff (Blossum-Henikoff, S and Henikoff, J.G., *PNAS*, 1992, **89**:10915-10919). The term "related kinases" also includes those containing residues with a similarity score of 80% or greater to the following ERK residues: I31, G37, A52, I103, E109, and N154. In a more preferred embodiment the similarity score is 85%, more preferably 90%, even more preferably 95%, 96%, 97% or 98%.

The present method is especially useful for treating a disease that is alleviated by the use of an inhibitor of ERK or related kinases. As used herein, unless otherwise indicated, the term "ERK" refers to all isoforms of the ERK enzyme including, but not limited to, ERK1, ERK2, ERK3, ERK4, ERK5, ERK6, and ERK7.

In an alternate embodiment, the methods of this invention that utilize compositions that do not contain an additional therapeutic agent, comprise the additional step of separately administering to said patient an additional therapeutic agent. When these additional therapeutic agents are administered separately they may be administered to the patient prior to, sequentially with or following administration of the compositions of this invention.

The compounds of this invention or pharmaceutically acceptable compositions thereof may also be incorporated into compositions for coating an implantable medical device, such as prostheses, artificial valves, vascular grafts, stents and catheters. Vascular stents, for example, have been used to overcome restenosis (re-narrowing of the vessel wall after injury). However, patients using stents or other

implantable devices risk clot formation or platelet activation. These unwanted effects may be prevented or mitigated by pre-coating the device with a pharmaceutically acceptable composition comprising a kinase inhibitor. Suitable coatings and the general preparation of coated implantable devices are described in US Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may be further covered by a suitable topcoat of fluorosilicone, polysaccharides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition. Implantable devices coated with a compound of this invention are another embodiment of the present invention.

In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

#### EXAMPLE 1

##### ERK Inhibition Assay

Compounds are assayed for the inhibition of ERK2 by a spectrophotometric coupled-enzyme assay (Fox et al (1998) *Protein Sci* 7, 2249). In this assay, a fixed concentration of activated ERK2 (10 nM) is incubated with various concentrations of the compound in DMSO (2.5 %) for 10 min. at 30°C in 0.1 M HEPES buffer, pH 7.5, containing 10 mM MgCl<sub>2</sub>, 2.5 mM phosphoenolpyruvate, 200 μM NADH, 150 μg/mL pyruvate kinase, 50 μg/mL lactate

dehydrogenase, and 200  $\mu$ M erktide peptide. The reaction is initiated by the addition of 65  $\mu$ M ATP. The rate of decrease of absorbance at 340 nm is monitored, which indicates the extent of uninhibited enzyme present in the assay. The  $IC_{50}$  is evaluated from the rate data as a function of inhibitor concentration.

## EXAMPLE 2

### ERK Inhibition Cell Proliferation Assay

Compounds may be assayed for the inhibition of ERK2 by a cell proliferation assay. In this assay, a complete media is prepared by adding 10% fetal bovine serum and penicillin/streptomycin solution to RPMI 1640 medium (JRH Biosciences). Colon cancer cells (HT-29 cell line) are added to each of 84 wells of a 96 well plate at a seeding density of 10,000 cells/well/150  $\mu$ L. The cells are allowed to attach to the plate by incubating at 37°C for 2 hours. A solution of test compound is prepared in complete media by serial dilution to obtain the following concentrations: 20  $\mu$ M, 6.7  $\mu$ M, 2.2  $\mu$ M, 0.74  $\mu$ M, 0.25  $\mu$ M, and 0.08  $\mu$ M. The test compound solution (50  $\mu$ L) is added to each of 72 cell-containing wells. To the 12 remaining cell-containing wells, only complete media (200  $\mu$ L) is added to form a control group in order to measure maximal proliferation. To the remaining 12 empty wells, complete media is added to form a vehicle control group in order to measure background. The plates are incubated at 37°C for 3 days. A stock solution of  $^3$ H-thymidine (1 mCi/mL, New England Nuclear, Boston, MA) is diluted to 20  $\mu$ Ci/mL in RPMI medium then 20  $\mu$ L of this solution is added to each well. The plates are further incubated at 37°C for 8 hours then harvested and analyzed for  $^3$ H-thymidine uptake using a liquid scintillation counter.


While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments, which utilize the compounds and methods of this invention.

- 5 Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments, which have been represented by way of example.



ClaimsWe Claim:

1. A compound of formula I:



or a pharmaceutically acceptable salt or derivative thereof, wherein:

- 5                   Ht is a heterocyclic ring selected from pyrrol-3-yl, [1,2,4]triazol-3-yl, [1,2,3]triazol-4-yl, or tetrazol-5-yl, said pyrrol-3-yl having R³ and Q<sub>n</sub>-R⁴ substituents, and said [1,2,4]triazol-3-yl or [1,2,3]triazol-4-yl substituted by either R³ or Q<sub>n</sub>-R⁴;
- 10                   T and Q are each independently selected from -C(R⁷)₂-, -C(O)-, -C(O)C(O)-, -C(O)NR⁷-, -C(O)NR⁷NR⁷-, -CO₂-, -OC(O)-, -NR⁷CO₂-, -O-, -NR⁷C(O)NR⁷-, -OC(O)NR⁷-, -NR⁷NR⁷-, -NR⁷C(O)-, -S-, -SO-, -SO₂-, -NR⁷-, -SO₂NR⁷-, or -NR⁷SO₂-, -NR⁷SO₂NR⁷-;
- 15                   m and n are each independently selected from zero or one;
- R² is selected from hydrogen, CN, halogen, R, N(R⁷)₂, OR, or OH;
- each R³ is independently selected from R⁷, F, Cl, -(CH₂)<sub>y</sub>N(R⁷)₂, -N(R⁷)₂, -OR⁷, -SR⁷, -NR⁷C(O)R⁷, -NR⁷C(O)N(R⁷)₂, -C(O)N(R⁷)₂, -SO₂R⁷, -NR⁷SO₂R⁷, -C(O)R⁷, -C(O)OR⁷, CN or -SO₂N(R⁷)₂;
- 20                   R⁴ is selected from -(CH₂)<sub>y</sub>R⁶, -(CH₂)<sub>y</sub>CH(R⁶)₂, -(CH₂)<sub>y</sub>CH(R⁸)CH(R⁶)₂, -N(R⁵)₂, -NR⁵(CH₂)<sub>y</sub>N(R⁵)₂;

each R is independently selected from an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, heteroaryl having 5-10 ring atoms, and heterocyclyl having 3-10 ring atoms;

5 each R<sup>5</sup> is independently selected from R, -R<sup>7</sup>, -C(O)R<sup>7</sup>, -CO<sub>2</sub>R, -C(O)N(R<sup>7</sup>)<sub>2</sub>, -SO<sub>2</sub>R<sup>7</sup>, -(CH<sub>2</sub>)<sub>y</sub>R<sup>6</sup>, or -(CH<sub>2</sub>)<sub>y</sub>CH(R<sup>6</sup>)<sub>2</sub>;

y is 0-6;

10 each R<sup>6</sup> is independently selected from R, -(CH<sub>2</sub>)<sub>y</sub>R, -OR, -CO<sub>2</sub>R, -(CH<sub>2</sub>)<sub>y</sub>N(R<sup>7</sup>)<sub>2</sub>, -N(R<sup>7</sup>)<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -NR<sup>7</sup>C(O)N(R<sup>7</sup>)<sub>2</sub>, -C(O)N(R<sup>7</sup>)<sub>2</sub>, -SO<sub>2</sub>R<sup>7</sup>, -NR<sup>7</sup>SO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -CN, -SO<sub>2</sub>N(R<sup>7</sup>)<sub>2</sub>;

15 each R<sup>7</sup> is independently selected from hydrogen or an optionally substituted C<sub>1-6</sub> aliphatic group, or two R<sup>7</sup> on the same nitrogen are taken together with the nitrogen to form a 5-8 membered ring heterocyclyl or heteroaryl ring;


each R<sup>8</sup> is independently selected from R, -(CH<sub>2</sub>)<sub>w</sub>OR<sup>7</sup>, -(CH<sub>2</sub>)<sub>w</sub>N(R<sup>5</sup>)<sub>2</sub>, or -(CH<sub>2</sub>)<sub>w</sub>SR<sup>7</sup>;

each w is independently 0-4;


provided that when Ht is tetrazol-5-yl, and m is 0, R<sup>2</sup> is other than H or an unsubstituted phenyl; when m is 0 and R<sup>2</sup>, R<sup>3</sup> and Q<sub>n</sub>-R<sup>4</sup> are all H, Ht is other than [1,2,3]triazole; and when T<sub>m</sub>-R<sup>2</sup> and Q<sub>n</sub>-R<sup>4</sup> are both C(O)OH, Ht is other than [1,2,3]triazol-5-yl.

2. The compound according to claim 1 wherein said compound is selected from the following:


-70-




II-A




II-B



II-C




II-D



II-E

or a pharmaceutically acceptable salt or derivative thereof.

3. The compound according to claim 2 wherein said compound is of formula II-A:



II-A

or a pharmaceutically acceptable salt or derivative thereof.

4. The compound according to any one of claims 1, 2, or 3 wherein said compound has one or more features selected from the group consisting of: (a)  $T_m R^2$  is hydrogen,  $N(R^7)_2$ , OH, 3-6 membered carbocyclyl, or an optionally substituted group selected from  $C_{1-6}$  aliphatic or a 5-6 membered aryl or heteroaryl ring; (b) Q is -CO-, -CO<sub>2</sub>-, -CONH-, -SO<sub>2</sub>-, -SO<sub>2</sub>NH-, -OC(O)NH-, -C(O)ONH-, or -CONHNH-; (c)  $R^4$  is  $-NR^5(CH_2)_yN(R^5)_2$ ,  $-(CH_2)_yR^6$ ,

$-(CH_2)_yCH(R^6)_2$ , or  $-(CH_2)_yCH(R^8)CH(R^6)_2$ ; (d)  $R^5$  is R,  $R^7$ , or  $-(CH_2)_yCH(R^6)_2$ ; and (e)  $R^6$  is an optionally substituted group selected from phenyl, 5-6 membered heteroaryl, or 5-6 membered heterocyclyl.

5

5. The compound according to claim 4 wherein:


(a)  $T_mR^2$  is hydrogen,  $N(R^7)_2$ , OH, 3-6 membered carbocyclyl, or an optionally substituted group selected from  $C_{1-6}$  aliphatic or a 5-6 membered aryl or heteroaryl ring; (b)  
10 Q is  $-CO-$ ,  $-CO_2-$ ,  $-CONH-$ ,  $-SO_2-$ ,  $-SO_2NH-$ ,  $-OC(O)NH-$ ,  $-C(O)ONH-$ , or  $-CONHNH-$ ; (c)  $R^4$  is  $-NR^5(CH_2)_yN(R^5)_2$ ,  $-(CH_2)_yR^6$ ,  $-(CH_2)_yCH(R^6)_2$ , or  $-(CH_2)_yCH(R^8)CH(R^6)_2$ ; (d)  $R^5$  is R,  $R^7$ , or  $-(CH_2)_yCH(R^6)_2$ ; and (e)  $R^6$  is an optionally substituted group selected from phenyl, 5-6 membered  
15 heteroaryl, or 5-6 membered heterocyclyl.

6. The compound according to claim 5 wherein:

(a)  $T_mR^2$  is selected from optionally substituted phenyl, methyl, ethyl, propyl, cyclopropyl, cyclohexyl,  $CH_2OCH_3$ ,  
20  $CH_2OH$ , OH,  $NH_2$ ,  $NHCH_3$ ,  $NHAc$ ,  $NHC(O)NHCH_3$ , or  $CH_2NHCH_3$ ; (b) Q is  $-CO-$ ,  $-CONH-$ ,  $-SO_2-$ , or  $-SO_2NH-$ ; (c)  $R^4$  is  $-(CH_2)_yR^6$ ,  $-(CH_2)_yCH(R^6)_2$ , or  $-(CH_2)_yCH(R^8)CH(R^6)_2$ , wherein  $R^8$  is OH or  $CH_2OH$ ; and (d)  $R^6$  is  $-CH_2OH$ ,  $-(CH_2)_2OH$ , isopropyl, or an optionally substituted group selected from pyrrolidin-1-yl,  
25 yl, morpholin-4-yl, piperidin-1-yl, piperazin-1-yl, 4-methyl[1,4]diazepan-1-yl, 4-phenyl-piperazine-1-yl, pyridin-3-yl, pyridin-4-yl, imidazolyl, furan-2-yl, 1,2,3,4-tetrahydroisoquinoline, tetrahydrofuran-2-yl, cyclohexyl, phenyl, or benzyl.

30


7. The compound according to claim 2 wherein said compound is selected from the following:



or a pharmaceutically acceptable salt or derivative thereof.

5

8. The compound according to claim 7 wherein said compound is of formula **III-A**:



**III-A**


10 or a pharmaceutically acceptable salt or derivative thereof.

9. The compound according to claim 7 wherein said compound has one or more features selected from the group consisting of: (a) when m is one, T is -NH- or -O-; (b) R<sup>2</sup> is an optionally substituted aliphatic or aryl group, more preferably an optionally substituted phenyl group; (c) R<sup>4</sup> is -N(R<sup>5</sup>)<sub>2</sub>, -NR<sup>5</sup>(CH<sub>2</sub>)<sub>y</sub>N(R<sup>5</sup>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>y</sub>R<sup>6</sup>, -(CH<sub>2</sub>)<sub>y</sub>CH(R<sup>6</sup>)<sub>2</sub>, or R<sup>6</sup>; (d) R<sup>5</sup> is R or R<sup>6</sup>; and/or (e) R<sup>6</sup> is an optionally substituted group selected from aryl, aralkyl, heteroaryl, heteroarylalkyl, heterocyclyl or heterocyclylalkyl group.

15

20

10. The compound according to claim 2 wherein said compound is selected from the following:



5 or a pharmaceutically acceptable salt or derivative thereof.

11. The compound according to claim 10 wherein said compound has one or more features selected from the group consisting of: (a)  $T_m R^2$  is hydrogen,  $N(R^7)_2$ , OH, 3-6 membered carbocyclyl, or an optionally substituted group selected from  $C_{1-6}$  aliphatic or a 5-6 membered aryl or heteroaryl ring; and (b)  $R^6$  is an optionally substituted 6-membered aryl, heteroaryl, or carbocyclic ring.

15


12. The compound according to claim 11 wherein: (a)  $T_m R^2$  is selected from optionally substituted phenyl, methyl, ethyl, propyl, cyclopropyl, cyclohexyl,  $CH_2OCH_3$ ,  $CH_2OH$ , OH,  $NH_2$ ,  $NHCH_3$ ,  $NHAc$ ,  $NHC(O)NHCH_3$ , or  $CH_2NHCH_3$ ; and (b)  $R^6$  is cyclohexyl or an optionally substituted phenyl or pyridyl ring.

20

13. The compound according to claim 2 wherein said compound is selected from the following:

25

-74-



II-A°,

II-B° or

II-D°

or a pharmaceutically acceptable salt or derivative thereof.


5

14. The compound according to claim 13 wherein said compound has one or more features selected from the group consisting of: (a) T<sub>m</sub>R<sup>2</sup> is hydrogen, N(R<sup>5</sup>)<sub>2</sub>, OH, 3-6 membered carbocyclyl, or an optionally substituted group selected from C<sub>1-6</sub> aliphatic or a 5-6 membered aryl or heteroaryl ring; and (b) R<sup>6</sup> is R or OR<sup>7</sup>, and R<sup>8</sup> is R<sup>7</sup> or OR<sup>7</sup>.

15. The compound according to claim 14 wherein: (a) T<sub>m</sub>R<sup>2</sup> is selected from optionally substituted phenyl, methyl, ethyl, propyl, cyclopropyl, cyclohexyl, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>OH, OH, NH<sub>2</sub>, NHCH<sub>3</sub>, NHAc, NHC(O)NHCH<sub>3</sub>, or CH<sub>2</sub>NHCH<sub>3</sub>; and (b) R<sup>6</sup> is OH, CH<sub>2</sub>OH, phenyl, pyridyl, or cyclohexyl, and R<sup>8</sup> is methyl, ethyl, OH, or CH<sub>2</sub>OH.

20



16. The compound according to claim 2 wherein said compound is of formula II-A, II-B or II-D










and is selected from any one of the following compounds, wherein  $R^3$  is H, m is zero, and n is one:




No.	$R^2$	$Q-R^4$
II-A-1, II-B-1, and II-C-1	phenyl	CON(Me) <sub>2</sub>
II-A-2, II-B-2, and II-C-2	phenyl	CO <sub>2</sub> Et
II-A-3, II-B-3, and II-C-3	3-NO <sub>2</sub> -phenyl	CONHNH <sub>2</sub>
II-A-4, II-B-4, and II-C-4	phenyl	CO(pyrrolidin-1-yl)
II-A-5, II-B-5, and II-C-5	phenyl	CONHCH <sub>2</sub> (Ph)
II-A-6, II-B-6, and II-C-6	3-NO <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-7, II-B-7, and II-C-7	4-Cl-phenyl	CO <sub>2</sub> Et
II-A-8, II-B-8, and II-C-8	4-OMe-phenyl	CO <sub>2</sub> Et
II-A-9, II-B-9, and II-C-9	3-NH <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-10, II-B-10, and II-C-10	3-OMe-phenyl	CO <sub>2</sub> Et
II-A-11, II-B-11, and II-C-11	4-F-phenyl	CO <sub>2</sub> Et
II-A-12, II-B-12, and II-C-12	4-NO <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-13, II-B-13, and II-C-13	3-Cl-phenyl	CO <sub>2</sub> Et
II-A-14, II-B-14, and II-C-14	3-F-phenyl	CO <sub>2</sub> Et
II-A-15, II-B-15, and II-C-15	phenyl	CO <sub>2</sub> H
II-A-16, II-B-16, and II-C-16	4-NH <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-17, II-B-17, and II-C-17	phenyl	CONHCH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>
II-A-18, II-B-18, and II-C-18	phenyl	CONHCH <sub>2</sub> (pyridin-3-yl)
II-A-19, II-B-19, and II-C-19	phenyl	CO(morpholin-4-yl)
II-A-20, II-B-20, and II-C-20	phenyl	CONH(isopropyl)
II-A-21, II-B-21, and II-C-21	phenyl	CO(4-Me-piperazin-1-yl)
II-A-22, II-B-22, and II-C-22	phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-23, II-B-23, and II-C-23	3-OMe-phenyl	CONMe <sub>2</sub>
II-A-24, II-B-24, and II-C-24	3-OMe-phenyl	CO(pyrrolidin-1-yl)
II-A-25, II-B-25, and II-C-25	3-OMe-phenyl	CONHCH <sub>2</sub> CH <sub>2</sub> N(Me) <sub>2</sub>
II-A-26, II-B-26, and II-C-26	3-OMe-phenyl	CONHCH <sub>2</sub> (pyridin-3-yl)
II-A-27, II-B-27, and II-C-27	3-OMe-phenyl	CO(morpholin-4-yl)
II-A-28, II-B-28, and II-C-28	3-OMe-phenyl	CONH(isopropyl)
II-A-29, II-B-29, and II-C-29	3-OMe-phenyl	CO(4-Me-














No.	R <sup>2</sup>	Q-R <sup>4</sup>
		piperazin-1-yl)
II-A-30, II-B-30, and II-C-30	3-OMe-phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-31, II-B-31, and II-C-31	4-NH <sub>2</sub> -phenyl	CO <sub>2</sub> Et
II-A-32, II-B-32, and II-C-32	H	CONMe <sub>2</sub>
II-A-33, II-B-33, and II-C-33	H	CO(pyrrolidin-1-yl)
II-A-34, II-B-34, and II-C-34	3-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-35, II-B-35, and II-C-35	4-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-36, II-B-36, and II-C-36	3-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-37, II-B-37, and II-C-37	4-(AcNH)-phenyl	CO <sub>2</sub> Et
II-A-38, II-B-38, and II-C-38	3-Cl-phenyl	CONHBn
II-A-39, II-B-39, and II-C-39	3,5-dichlorophenyl	
II-A-40, II-B-40, and II-C-40	3-Br-phenyl	CONH(3,4-difluorophenyl)
II-A-41, II-B-41, and II-C-41	3-Cl-phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-42, II-B-42, and II-C-42	4-OH-3-I-5-nitrophenyl	CONH(2-OH-1-Ph-ethyl)
II-A-43, II-B-43, and II-C-43	3-Br-phenyl	
II-A-44, II-B-44, and II-C-44	5-NH <sub>2</sub> -4-OH-3-I-phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-45, II-B-45, and II-C-45	3-Br-phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-46, II-B-46, and II-C-46	3-Br-phenyl	CONHCH <sub>2</sub> (3-MeO-phenyl)
II-A-47, II-B-47, and II-C-47	3-Br-phenyl	CONHCH <sub>2</sub> (3-CF <sub>3</sub> -phenyl)
II-A-48, II-B-48, and II-C-48	3,5-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-49, II-B-49, and II-C-49	3-CF <sub>3</sub> -phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-50, II-B-50, and II-C-50	3-Cl-phenyl	CONHCH <sub>2</sub> Ph
II-A-51, II-B-51, and II-C-51	3,5-dichlorophenyl	CONHOCH <sub>2</sub> Ph
II-A-52, II-B-52, and II-C-52	4-OH-3-I-5-nitrophenyl	CONHCH <sub>2</sub> Ph
II-A-53, II-B-53, and II-C-53	3-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)



No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-54, II-B-54, and II-C-54	3,4-dichlorophenyl	CONHOCH <sub>2</sub> Ph
II-A-55, II-B-55, and II-C-55	3-Br-phenyl	CONHCH <sub>2</sub> (4-SO <sub>2</sub> Me-phenyl)
II-A-56, II-B-56, and II-C-56	3-Br-phenyl	CONHNH(3-CF <sub>3</sub> -phenyl)
II-A-57, II-B-57, and II-C-57	3-Cl-phenyl	CONHOCH <sub>2</sub> Ph
II-A-58, II-B-58, and II-C-58	3-Br-phenyl	
II-A-59, II-B-59, and II-C-59	3-Br-phenyl	
II-A-60, II-B-60, and II-C-60	3-Br-phenyl	CONHCH <sub>2</sub> (2-Me-phenyl)
II-A-61, II-B-61, and II-C-61	3,4-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-62, II-B-62, and II-C-62	3-Br-phenyl	CONH(1-Ph-propyl)
II-A-63, II-B-63, and II-C-63	3-F-phenyl	CONHCH <sub>2</sub> Ph
II-A-64, II-B-64, and II-C-64	3,4-dichlorophenyl	
II-A-65, II-B-65, and II-C-65	3-Br-phenyl	
II-A-66, II-B-66, and II-C-66	3,5-dichlorophenyl	CON(Me)(Et)
II-A-67, II-B-67, and II-C-67	3-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-68, II-B-68, and II-C-68	3-Br-phenyl	CONHCH <sub>2</sub> (3,5-dimethoxyphenyl)
II-A-69, II-B-69, and II-C-69	3-Br-phenyl	CONHCH <sub>2</sub> (2-OMe-phenyl)
II-A-70, II-B-70, and II-C-70	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-71, II-B-71, and II-C-71	3-Cl-4-F-phenyl	CON(Me)(Et)
II-A-72, II-B-72, and II-C-72	3-Br-phenyl	CONH(2-OH-1-Ph-ethyl)
II-A-73, II-B-73, and II-C-73	3-NH <sub>2</sub> -phenyl	CONHCH <sub>2</sub> Ph
II-A-74, II-B-74, and II-C-74	3,4-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-75, II-B-75, and II-C-75	3-Me-phenyl	CONH(2-OH-1-Ph-

No.	R <sup>2</sup>	Q-R <sup>4</sup>
		ethyl)
II-A-76, II-B-76, and II-C-76	3,5-dichlorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-77, II-B-77, and II-C-77	3-Cl-4-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-78, II-B-78, and II-C-78	3,5-dichlorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-79, II-B-79, and II-C-79	3-NO <sub>2</sub> -phenyl	CONHCH <sub>2</sub> Ph
II-A-80, II-B-80, and II-C-80	3-F-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-81, II-B-81, and II-C-81	3-Cl-2-F-phenyl	CON (Me) (Et)
II-A-82, II-B-82, and II-C-82	3-Cl-2-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-83, II-B-83, and II-C-83	3-Br-phenyl	
II-A-84, II-B-84, and II-C-84	3-Cl-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-85, II-B-85, and II-C-85	3,4-difluorophenyl	CONHOCH <sub>2</sub> Ph
II-A-86, II-B-86, and II-C-86	3-Br-phenyl	CONH (3-OH-1-Ph-propyl)
II-A-87, II-B-87, and II-C-87	3-Br-phenyl	
II-A-88, II-B-88, and II-C-88	3,4-difluorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-89, II-B-89, and II-C-89	3-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-90, II-B-90, and II-C-90	3-Me-phenyl	CONHCH <sub>2</sub> Ph
II-A-91, II-B-91, and II-C-91	3-Br-phenyl	
II-A-92, II-B-92, and II-C-92	4-Cl-phenyl	CONHCH <sub>2</sub> Ph
II-A-93, II-B-93, and II-C-93	3-Cl-phenyl	CON (Me) (Et)
II-A-94, II-B-94, and II-C-94	3-Br-phenyl	CONHCH <sub>2</sub> (4-SO <sub>2</sub> NH <sub>2</sub> -phenyl)
II-A-95, II-B-95, and II-C-95	3-OH-phenyl	CONHCH <sub>2</sub> Ph
II-A-96, II-B-96, and II-C-96	3-Me-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-97, II-B-97, and II-C-97	Phenyl	CONHCH <sub>2</sub> Ph
II-A-99, II-B-98, and II-C-98	2,5-difluorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-99, II-B-99, and II-C-99	4-Cl-phenyl	CONHOCH <sub>2</sub> Ph
II-A-100, II-B-100, and II-C-100	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)

No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-101, II-B-101, and II-C-101	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-102, II-B-102, and II-C-102	3-Br-phenyl	 CO(4-OH-4-Ph-piperidin-1-yl)
II-A-103, II-B-103, and II-C-103	2,3-difluorophenyl	CONHOCH <sub>2</sub> Ph
II-A-104, II-B-104, and II-C-104	3-Cl-phenyl	CO(morpholin-4-yl)
II-A-105, II-B-105, and II-C-105	3-Br-phenyl	
II-A-106, II-B-106, and II-C-106	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-107, II-B-107, and II-C-107	3-Cl-4-F-phenyl	CO(morpholin-4-yl)
II-A-108, II-B-108, and II-C-108	3-Cl-4-F-phenyl	CON(Me)(Et)
II-A-109, II-B-109, and II-C-109	3-Br-phenyl	CONHCH <sub>2</sub> (4-NH <sub>2</sub> -phenyl)
II-A-110, II-B-110, and II-C-110	3-Br-phenyl	
II-A-111, II-B-111, and II-C-111	4-F-phenyl	CONHCH <sub>2</sub> Ph
II-A-112, II-B-112, and II-C-112	3,5-dichlorophenyl	CO(morpholin-4-yl)
II-A-113, II-B-113, and II-C-113	2,5-difluorophenyl	CONHOCH <sub>2</sub> Ph
II-A-114, II-B-114, and II-C-114	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-115, II-B-115, and II-C-115	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-116, II-B-116, and II-C-116	3,4-difluorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-117, II-B-117, and II-C-117	4-OMe-phenyl	CONHCH <sub>2</sub> Ph
II-A-118, II-B-118, and II-C-118	3-Br-phenyl	CONHCH <sub>2</sub> (2,4,6-trimethoxyphenyl)
II-A-119, II-B-119, and II-C-119	3-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-120, II-B-120, and II-C-120	3,4-difluorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)

No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-121, II-B-121, and II-C-121	3-Cl-2-F-phenyl	
II-A-122, II-B-122, and II-C-122	3-Br-phenyl	
II-A-123, II-B-123, and II-C-123	3-Br-phenyl	
II-A-124, II-B-124, and II-C-124	3-Br-phenyl	CONHCH <sub>2</sub> (2,5-dimethoxyphenyl)
II-A-125, II-B-125, and II-C-125	3,5-dichlorophenyl	
II-A-126, II-B-126, and II-C-126	3-Br-phenyl	
II-A-127, II-B-127, and II-C-127	3,4-dichlorophenyl	CO (morpholin-4-yl)
II-A-128, II-B-128, and II-C-128	3-Br-phenyl	
II-A-129, II-B-129, and II-C-129	3-Cl-2-F-phenyl	CO (morpholin-4-yl)
II-A-130, II-B-130, and II-C-130	3-Br-phenyl	CONHCH <sub>2</sub> CH <sub>2</sub> OH
II-A-131, II-B-131, and II-C-131	3-NH <sub>2</sub> -phenyl	CONHCH <sub>2</sub> Ph
II-A-132, II-B-132, and II-C-132	3-MeOC(O)-phenyl	CONHCH <sub>2</sub> Ph
II-A-133, II-B-133, and II-C-133	4-MeO-phenyl	CONHOCH <sub>2</sub> Ph
II-A-134, II-B-134, and II-C-134	Phenyl	CO (pyrrolidin-1-yl)
II-A-135, II-B-135, and II-C-135	3-MeO-phenyl	CO (morpholin-4-yl)
II-A-136, II-B-136, and II-C-136	3-Cl-phenyl	CO (4-Me-piperidin-1-yl)
II-A-137, II-B-137, and II-C-137	3-NO <sub>2</sub> -phenyl	CONH <sub>2</sub> NH <sub>2</sub>
II-A-138, II-B-138, and II-C-138	3-Br-phenyl	
II-A-139, II-B-139, and II-C-139	3-Br-phenyl	
II-A-140, II-B-140, and II-C-140	3-Cl-phenyl	CONHPh













No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-141, II-B-141, and II-C-141	2,3-difluorophenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-142, II-B-142, and II-C-142	3-Cl-phenyl	
II-A-143, II-B-143, and II-C-143	Phenyl	CON(Me) <sub>2</sub>
II-A-144, II-B-144, and II-C-144	3-OMe-phenyl	CO(pyrrolidin-1-yl)
II-A-145, II-B-145, and II-C-145	3-OMe-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-146, II-B-146, and II-C-146	4-F-phenyl	CONHOCH <sub>2</sub> Ph
II-A-147, II-B-147, and II-C-147	3-OMe-phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-148, II-B-148, and II-C-148	3-NO <sub>2</sub> -phenyl	COOEt
II-A-149, II-B-149, and II-C-149	Phenyl	CONHCH <sub>2</sub> (furan-2-yl)
II-A-150, II-B-150, and II-C-150	Phenyl	CO(morpholin-4-yl)
II-A-151, II-B-151, and II-C-151	3-Cl-phenyl	COOEt
II-A-152, II-B-152, and II-C-152	3-Br-phenyl	CONHMe
II-A-153, II-B-153, and II-C-153	Phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-154, II-B-154, and II-C-154	3-OMe-phenyl	CON(Me) <sub>2</sub>
II-A-155, II-B-155, and II-C-155	3-Cl-phenyl	
II-A-156, II-B-156, and II-C-156	3-Br-phenyl	
II-A-157, II-B-157, and II-C-157	3-Br-phenyl	COOEt
II-A-158, II-B-158, and II-C-158	phenyl	CONH(iPr)
II-A-159, II-B-159, and II-C-159	3-OMe-phenyl	CONH(iPr)
II-A-160, II-B-160, and II-C-160	3-COOH-phenyl	CONH(iPr)
II-A-161, II-B-161, and II-C-161	3-Br-phenyl	CONHO(iPr)
II-A-162, II-B-162, and II-C-162	3-F-phenyl	COOEt
II-A-163, II-B-163, and II-C-163	3-OMe-phenyl	CO(4-Me-piperidin-1-yl)
II-A-164, II-B-164, and II-C-164	4-NH <sub>2</sub> -phenyl	COOEt
II-A-165, II-B-165, and II-C-165	4-NO <sub>2</sub> -phenyl	COOEt
II-A-166, II-B-166, and II-C-166	phenyl	CO(4-Me-piperidin-1-yl)

No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-167, II-B-167, and II-C-167	4-Cl-phenyl	COOEt
II-A-168, II-B-168, and II-C-168	4-OMe-phenyl	COOEt
II-A-169, II-B-169, and II-C-169	phenyl	COOEt
II-A-170, II-B-170, and II-C-170	3-OMe-phenyl	COOEt
II-A-171, II-B-171, and II-C-171	4-F-phenyl	COOEt
II-A-172, II-B-172, and II-C-172	3-NH <sub>2</sub> -phenyl	COOEt
II-A-173, II-B-173, and II-C-173	3-Cl-phenyl	COOH
II-A-174, II-B-174, and II-C-174	3-Cl-phenyl	
II-A-175, II-B-175, and II-C-175	3-Cl-phenyl	
II-A-176, II-B-176, and II-C-176	3-OMe-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-177, II-B-177, and II-C-177	3,5-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-178, II-B-178, and II-C-178	4-F-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-179, II-B-179, and II-C-179	4-OMe-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-180, II-B-180, and II-C-180	2,5-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-181, II-B-181, and II-C-181	2,5-difluorophenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-182, II-B-182, and II-C-182	4-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-183, II-B-183, and II-C-183	4-OMe-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-184, II-B-184, and II-C-184	3-F-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-185, II-B-185, and II-C-185	3-OMe-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-186, II-B-186, and II-C-186	2,5-dimethoxyphenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-187, II-B-187, and II-C-187	2,3-difluorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-188, II-B-188, and II-C-188	2,5-difluorophenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-189, II-B-189, and II-C-189	4-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-190, II-B-190, and II-C-190	4-OMe-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-191, II-B-191, and II-C-191	3-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-192, II-B-192, and II-C-192	3-OMe-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-193, II-B-193, and II-C-193	2,5-dimethoxyphenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-194, II-B-194, and II-C-194	3,4-difluorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-195, II-B-195, and II-C-195	2,3-difluorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-196, II-B-196, and II-C-196	2,5-difluorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-197, II-B-197, and II-C-197	4-F-phenyl	CO(morpholin-4-yl)
II-A-198, II-B-198, and II-C-198	4-OMe-phenyl	CO(morpholin-4-yl)
II-A-199, II-B-199, and II-C-199	3-F-phenyl	CO(morpholin-4-yl)
II-A-200, II-B-200, and II-C-200	2,5-dimethoxyphenyl	CO(morpholin-4-yl)
II-A-201, II-B-201, and II-C-201	3,4-difluorophenyl	CO(morpholin-4-yl)
II-A-202, II-B-202, and II-C-202	2,3-difluorophenyl	CO(morpholin-4-yl)
II-A-203, II-B-203, and II-C-203	2,5-difluorophenyl	CO(morpholin-4-yl)
II-A-204, II-B-204, and II-C-204	4-F-phenyl	CO(4-Me-piperidin-1-yl)
II-A-205, II-B-205, and II-C-205	4-OMe-phenyl	CO(4-Me-piperidin-1-yl)
II-A-206, II-B-206, and II-C-206	3-F-phenyl	CO(4-Me-piperidin-1-yl)
II-A-207, II-B-207, and II-C-207	2,5-dimethoxyphenyl	CO(4-Me-piperidin-1-yl)
II-A-208, II-B-208, and II-C-208	3,4-difluorophenyl	CO(4-Me-piperidin-1-yl)
II-A-209, II-B-209, and II-C-209	2,3-difluorophenyl	CO(4-Me-piperidin-1-yl)
II-A-210, II-B-210, and II-C-210	2,5-difluorophenyl	CO(4-Me-piperidin-1-yl)
II-A-211, II-B-211, and II-C-211	4-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-212, II-B-212, and II-C-212	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-213, II-B-213, and II-C-213	benzo[1,3]dioxo-5-yl	CONHCH <sub>2</sub> (pyrid-4-yl)



No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-214, II-B-214, and II-C-214	4-Cl-phenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-215, II-B-215, and II-C-215	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-216, II-B-216, and II-C-216	benzo[1,3]dioxo-5-yl	CONHCH <sub>2</sub> (pyrid-3-yl)
II-A-217, II-B-217, and II-C-217	4-Cl-phenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-218, II-B-218, and II-C-218	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-219, II-B-219, and II-C-219	benzo[1,3]dioxo-5-yl	CONHCH <sub>2</sub> (tetrahydrofuran-2-yl)
II-A-220, II-B-220, and II-C-220	4-Cl-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-221, II-B-221, and II-C-221	3,4-dichlorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-222, II-B-222, and II-C-222	3-Cl-2-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-223, II-B-223, and II-C-223	3-Cl-4-F-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-224, II-B-224, and II-C-224	3,4-dimethoxyphenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-225, II-B-225, and II-C-225	benzo[1,3]dioxo-5-yl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-226, II-B-226, and II-C-226	3,5-dichlorophenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-227, II-B-227, and II-C-227	4-Cl-phenyl	CO (morpholin-4-yl)
II-A-228, II-B-228, and II-C-228	3,4-dimethoxyphenyl	CO (morpholin-4-yl)
II-A-229, II-B-229, and II-C-229	benzo[1,3]dioxo-5-yl	CO (morpholin-4-yl)
II-A-230, II-B-230, and II-C-230	4-Cl-phenyl	CO (4-Me-piperidin-1-yl)
II-A-231, II-B-231, and II-C-231	3,4-dichlorophenyl	CO (4-Me-piperidin-1-yl)
II-A-232, II-B-232, and II-C-232	3-Cl-2-F-phenyl	CO (4-Me-piperidin-1-yl)
II-A-233, II-B-233, and II-C-233	3-Cl-4-F-phenyl	CO (4-Me-piperidin-1-yl)
II-A-234, II-B-234, and II-C-234	3,4-dimethoxyphenyl	CO (4-Me-piperidin-1-yl)
II-A-235, II-B-235, and II-C-235	benzo[1,3]dioxo-5-yl	CO (4-Me-piperidin-1-yl)
II-A-236, II-B-236, and II-C-236	3,5-dichlorophenyl	CO (4-Me-piperidin-1-yl)

No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-237, II-B-237, and II-C-237	2,3-difluorophenyl	CON(Me) (Et)
II-A-238, II-B-238, and II-C-238	4-F-phenyl	
II-A-239, II-B-239, and II-C-239	3-OMe-phenyl	
II-A-240, II-B-240, and II-C-240	2,5-dimethoxyphenyl	
II-A-241, II-B-241, and II-C-241	3,4-difluorophenyl	
II-A-242, II-B-242, and II-C-242	2,3-difluorophenyl	
II-A-243, II-B-243, and II-C-243	2,5-difluorophenyl	
II-A-244, II-B-244, and II-C-244	3-MeO-phenyl	CONHOCH <sub>2</sub> Ph
II-A-245, II-B-245, and II-C-245	2,5-dimethoxyphenyl	CONHOCH <sub>2</sub> Ph
II-A-246, II-B-246, and II-C-246	3-F-phenyl	
II-A-247, II-B-247, and II-C-247	3-MeO-phenyl	
II-A-248, II-B-248, and II-C-248	3,4-difluorophenyl	
II-A-249, II-B-249, and II-C-249	2,3-difluorophenyl	
II-A-250, II-B-250, and II-C-250	3-Cl-phenyl	
II-A-251, II-B-251, and II-C-251	4-Cl-phenyl	













No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-252, II-B-252, and II-C-252	4-Cl-phenyl	
II-A-253, II-B-253, and II-C-253	3,4-dichlorophenyl	
II-A-254, II-B-254, and II-C-254	3,4-dichlorophenyl	
II-A-255, II-B-255, and II-C-255	3-Cl-2-F-phenyl	
II-A-256, II-B-256, and II-C-256	3-Cl-4-F-phenyl	
II-A-257, II-B-257, and II-C-257	3-Cl-4-F-phenyl	
II-A-258, II-B-258, and II-C-258	3,4-dimethoxyphenyl	CON(Me) (Et)
II-A-259, II-B-259, and II-C-259	3,4-dimethoxyphenyl	
II-A-260, II-B-260, and II-C-260	3,4-dimethoxyphenyl	CONHOCH <sub>2</sub> Ph
II-A-261, II-B-261, and II-C-261	3,4-dimethoxyphenyl	
II-A-262, II-B-262, and II-C-262	benzo[1,3]dioxo-5-yl	CON(Me) (Et)
II-A-263, II-B-263, and II-C-263	benzo[1,3]dioxo-5-yl	
II-A-264, II-B-264, and II-C-264	benzo[1,3]dioxo-5-yl	CONHOCH <sub>2</sub> Ph
II-A-265, II-B-265, and II-C-265	benzo[1,3]dioxo-5-yl	
II-A-266, II-B-266, and II-C-266	3,5-dichlorophenyl	
II-A-267, II-B-267, and II-C-267	3-Br-phenyl	

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-268, II-B-268, and II-C-268	3-Br-phenyl	
II-A-269, II-B-269, and II-C-269	3-Br-phenyl	
II-A-270, II-B-270, and II-C-270	3-Br-phenyl	
II-A-271, II-B-271, and II-C-271	3-Br-phenyl	
II-A-272, II-B-272, and II-C-272	3-Br-phenyl	
II-A-273, II-B-273, and II-C-273	3-Br-phenyl	
II-A-274, II-B-274, and II-C-274	3-Br-phenyl	
II-A-275, II-B-275, and II-C-275	3-Br-phenyl	
II-A-276, II-B-276, and II-C-276	3-Br-phenyl	
II-A-277, II-B-277, and II-C-277	3-Br-phenyl	
II-A-278, II-B-278, and II-C-278	3-Br-phenyl	
II-A-279, II-B-279, and II-C-279	3-Br-phenyl	
II-A-280, II-B-280, and II-C-280	3-Br-phenyl	CONH(CH <sub>2</sub> ) <sub>2</sub> COOH
II-A-281, II-B-281, and II-C-281	3-Br-phenyl	
II-A-282, II-B-282, and II-C-282	3-Br-phenyl	CONHCH <sub>2</sub> (4-COOH-

No.	R <sup>2</sup>	Q-R <sup>4</sup>
		phenyl)
II-A-283, II-B-283, and II-C-283	3-Br-phenyl	
II-A-284, II-B-284, and II-C-284	3-Br-phenyl	
II-A-285, II-B-285, and II-C-285	3-NO <sub>2</sub> -phenyl	CONHCH <sub>2</sub> phenyl
II-A-286, II-B-286, and II-C-286	3-Cl-phenyl	CONHCH <sub>2</sub> (1-Et-pyrrolidin-2-yl)
II-A-287, II-B-287, and II-C-287	3-(N-Et-NHCO)-phenyl	CONHCH <sub>2</sub> phenyl
II-A-288, II-B-288, and II-C-288	3-Br-phenyl	
II-A-289, II-B-289, and II-C-289	3-NO <sub>2</sub> -phenyl	CONHCH <sub>2</sub> (pyrid-4-yl)
II-A-290, II-B-290, and II-C-290	3-Br-phenyl	
II-A-291, II-B-291, and II-C-291	3-F-phenyl	CON(Me)(Et)
II-A-292, II-B-292, and II-C-292	3-MeO-phenyl	CON(Me)(Et)
II-A-293, II-B-293, and II-C-293	3-Br-phenyl	
II-A-294, II-B-294, and II-C-294	3-Br-phenyl	
II-A-295, II-B-295, and II-C-295	3-Br-phenyl	

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-296, II-B-296, and II-C-296	3-Br-phenyl	
II-A-297, II-B-297, and II-C-297	phenyl	CONH(CH <sub>2</sub> ) <sub>2</sub> NMe <sub>2</sub>
II-A-298, II-B-298, and II-C-298	3-MeO-phenyl	CONH(CH <sub>2</sub> ) <sub>2</sub> NMe <sub>2</sub>
II-A-299, II-B-299, and II-C-299	3-Br-phenyl	CONHCH <sub>2</sub> phenyl
II-A-300, II-B-300, and II-C-300	3-Cl-phenyl	
II-A-301, II-B-301, and II-C-301	3-Cl-phenyl	
II-A-302, II-B-302, and II-C-302	3-Cl-phenyl	
II-A-303, II-B-303, and II-C-303	3-Cl-phenyl	
II-A-304, II-B-304, and II-C-304	3-Cl-phenyl	
II-A-305, II-B-305, and II-C-305	3-Cl-phenyl	
II-A-306, II-B-306, and II-C-306	3-Cl-phenyl	
II-A-307, II-B-307, and II-C-307	3-Cl-phenyl	
II-A-308, II-B-308, and II-C-308	3-Cl-phenyl	
II-A-309, II-B-309, and II-C-309	3-Cl-phenyl	

No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-310, II-B-310, and II-C-310	3,5-dichlorophenyl	
II-A-311, II-B-311, and II-C-311	3-Br-5-CF <sub>3</sub> -phenyl	
II-A-312, II-B-312, and II-C-312	3-Cl-phenyl	
II-A-313, II-B-313, and II-C-313	3,5-dichlorophenyl	
II-A-314, II-B-314, and II-C-314	3-Cl-4-CN-phenyl	
II-A-v15, II-B-315, and II-C-315	3-Cl-4-CH <sub>2</sub> OH-phenyl	
II-A-316, II-B-316, and II-C-316	3-Cl-4-CH <sub>2</sub> NH <sub>2</sub> -phenyl	
II-A-317, II-B-317, and II-C-317		
II-A-318, II-B-318, and II-C-318		
II-A-319, II-B-319, and II-C-319		
II-A-320, II-B-320, and II-C-320		


No .	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-321, II-B-321, and II-C-321		
II-A-322, II-B-322, and II-C-322		
II-A-323, II-B-323, and II-C-323	CH <sub>2</sub> Ph	CON(Me) <sub>2</sub>
II-A-324, II-B-324, and II-C-324	cyclopentylmethyl	CO <sub>2</sub> NHCH <sub>2</sub> Ph
II-A-325, II-B-325, and II-C-325	isopropyl	CN
II-A-326, II-B-326, and II-C-326	3-Cl-phenyl	NHCOCH <sub>2</sub> Ph
II-A-327, II-B-327, and II-C-327	3-Cl-phenyl	NHSO <sub>2</sub> -morpholin-4-yl
II-A-328, II-B-328, and II-C-328	3-Cl-phenyl	NHCONHCH <sub>2</sub> Ph
II-A-329, II-B-329, and II-C-329	3-Cl-phenyl	NHCO <sub>2</sub> -tetrahydrofuran-2-yl
II-A-330, II-B-330, and II-C-330	CH <sub>2</sub> Ph	CONHCH <sub>2</sub> Ph
II-A-331, II-B-331, and II-C-331	Me	CONHCH <sub>2</sub> Ph
II-A-332, II-B-332, and II-C-332	isopropyl	CONHCH <sub>2</sub> Ph
II-A-333, II-B-333, and II-C-333	H	CON(Me) <sub>2</sub>
II-A-334, II-B-334, and II-C-334		
II-A-335, II-B-335, and II-C-335		
II-A-336, II-B-336, and II-C-336		
II-A-337, II-B-337, and II-C-337		



No.	R <sup>2</sup>	Q-R <sup>4</sup>
II-A-338, II-B-338, and II-C-338		
II-A-339, II-B-339, and II-C-339		
II-A-340, II-B-340, and II-C-340		
II-A-341, II-B-341, and II-C-341		














17. The compound according to claim 16 wherein said compound is of formula II-A.















5 18. The compound according to claim 2 wherein said compound is of formula II-A, II-B or II-D














and is selected from any one of the following compounds:

No.	T <sub>m</sub> -R <sup>2</sup>	Q <sub>n</sub> -R <sup>4</sup>
II-A-342, II-B-342, and II-C-342	2-F-3-Cl-phenyl	

No.	T <sub>m</sub> -R <sup>2</sup>	Q <sub>n</sub> -R <sup>4</sup>
II-A-343, II-B-343, and II-C-343	methyl	
II-A-344, II-B-344, and II-C-344	methyl	
II-A-345, II-B-345, and II-C-345	Methyl	
II-A-346, II-B-346, and II-C-346	3,5-dichlorophenyl	
II-A-347, II-B-347, and II-C-347	3-F,5-CF <sub>3</sub> -phenyl	
II-A-348, II-B-348, and II-C-348	Methyl	
II-A-349, II-B-349, and II-C-349	H	
II-A-350, II-B-350, and II-C-350	Methyl	
II-A-351, II-B-351, and II-C-351	Methyl	
II-A-352, II-B-352, and II-C-352	Methyl	
II-A-353, II-B-353, and II-C-353	Cyclohexyl	
II-A-354, II-B-354, and II-C-354	Cyclopropyl	
II-A-355, II-B-355, and II-C-355	Methyl	







No.	$T_m-R^2$	$Q_n-R^4$
II-A-356, II-B-356, and II-C-356	Methyl	
II-A-357, II-B-357, and II-C-357	$CH_2OCH_3$	
II-A-358, II-B-358, and II-C-358	$CH_2OH$	
II-A-359, II-B-359, and II-C-359	Methyl	
II-A-360, II-B-360, and II-C-360	Methyl	
II-A-361, II-B-361, and II-C-361	Methyl	
II-A-362, II-B-362, and II-C-362	Methyl	
II-A-363, II-B-363, and II-C-363	Methyl	
II-A-364, II-B-364, and II-C-364	H	
II-A-365, II-B-365, and II-C-365	H	
II-A-366, II-B-366, and II-C-366	H	
II-A-367, II-B-367, and II-C-367	H	
II-A-368, II-B-368, and II-C-368	H	
II-A-369, II-B-369, and II-C-369	H	







No.	T <sub>m</sub> -R <sup>2</sup>	Q <sub>n</sub> -R <sup>4</sup>
II-A-370, II-B-370, and II-C-370	Methyl	
II-A-371, II-B-371, and II-C-371	Methyl	
II-A-372, II-B-372, and II-C-372	Methyl	
II-A-373, II-B-373, and II-C-373	Methyl	
II-A-374, II-B-374, and II-C-374	Methyl	
II-A-375, II-B-375, and II-C-375	Methyl	
II-A-376, II-B-376, and II-C-376	Methyl	
II-A-377, II-B-377, and II-C-377	Methyl	
II-A-378, II-B-378, and II-C-378	Methyl	
II-A-379, II-B-379, and II-C-379	Methyl	
II-A-380, II-B-380, and II-C-380	methyl	







No.	T <sub>m</sub> -R <sup>2</sup>	Q <sub>n</sub> -R <sup>4</sup>
II-A-381, II-B-381, and II-C-381	H	
II-A-382, II-B-382, and II-C-382	H	
II-A-383, II-B-383, and II-C-383		
II-A-384, II-B-384, and II-C-384		
II-A-385, II-B-385, and II-C-385		
II-A-386, II-B-386, and II-C-386		
II-A-387, II-B-387, and II-C-387		
II-A-388, II-B-388, and II-C-388		
II-A-389, II-B-389, and II-C-389		
II-A-390, II-B-390, and II-C-390		

19. The compound according to claim 18 wherein said compound is of formula II-A.







20. The compound according to claim 2 wherein said compound is selected from any one of the following compounds:




No.	Structure
II-A 1	
II-A 2	
II-A 3	
II-A 4	
II-A 5	
II-A 6	

No.	Structure
II-A 7	
II-A 8	
II-B 1	
II-B 2	
II-B 3	
II-B 4	



No.	Structure
II-B 5	
II-B 6	
II-C 1	
II-C 2	
II-C 3	
II-C 4	

















No.	Structure
II-C 5	
II-C 6	
II-C 7	
II-E 1	
II-E 2	
II-E 3	








No.	Structure
II-E 4	
II-E 5	
II-E 6	








21. The compound according to claim 2 wherein said compound is selected from any of the following compounds:

No.	Structure
II-A 9	
II-A 10	

No.	Structure
II-A 11	
II-A 12	
II-A 13	
II-A 14	
II-A 15	
II-A 16	
II-A 17	

No.	Structure
II-A 18	
II-A 19	
II-A 20	
II-A 21	
II-A 22	
II-A 23	
II-A 24	

No.	Structure
II-A 25	
II-A 26	
II-A 27	
II-A 28	
II-A 29	
II-A 30	
II-A 31	

No.	Structure
II-A 32	
II-A 33	
II-A 34	
II-A 35	
II-A 36	
II-A 37	
II-A 38	

22. A composition comprising a compound according to claim 1 in an amount to detectably inhibit ERK kinase activity, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

5

23. The composition according to claim 22, additionally comprising an additional therapeutic agent selected from an anti-proliferative agent, an anti-inflammatory agent, an immunomodulatory agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating liver disease, an anti-viral agent, an agent for treating blood disorders, an agent for treating diabetes, or an agent for treating immunodeficiency disorders.

10  
15

24. A method of inhibiting ERK kinase activity in a biological sample comprising the step of contacting said biological sample with:

- a) a compound according to claim 1; or
- b) a composition according to claim 22.

20

25. A method of treating or lessening the severity of an ERK-mediated disease or condition in a patient comprising the step of administering to said patient a composition according to claim 22.

25

26. A method of treating or lessening the severity of an cancer, stroke, diabetes, hepatomegaly, cardiovascular disease, Alzheimer's disease, cystic fibrosis, viral disease, autoimmune diseases, atherosclerosis, restenosis, psoriasis, allergic disorders, inflammation, neurological disorders, a hormone-related disease, conditions associated with organ transplantation, immunodeficiency disorders, destructive

30

bone disorders, proliferative disorders, infectious diseases, conditions associated with cell death, thrombin-induced platelet aggregation, chronic myelogenous leukemia (CML), liver disease, pathologic  
5 immune conditions involving T cell activation, or CNS disorders, comprising the step of administering to said patient a composition according to claim 22.

27. The method according to claim 26, wherein  
10 said method is used to treat or prevent cancer.

28. The method according to claim 27, wherein said method is used to treat or prevent a cancer selected from breast; ovary; cervix; prostate; testis,  
15 genitourinary tract; esophagus; larynx, glioblastoma; neuroblastoma; stomach; skin, keratoacanthoma; lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma; bone; colon, adenoma; pancreas, adenocarcinoma; thyroid, follicular carcinoma,  
20 undifferentiated carcinoma, papillary carcinoma; seminoma; melanoma; sarcoma; bladder carcinoma; liver carcinoma and biliary passages; kidney carcinoma; myeloid disorders; lymphoid disorders, Hodgkin's, hairy cells; buccal cavity and pharynx (oral), lip, tongue, mouth,  
25 pharynx; small intestine; colon-rectum, large intestine, rectum; brain and central nervous system; or leukemia.

29. The method according to claim 26, wherein said method is used to treat or prevent cardiovascular  
30 disease.

30. The method according to claim 29, wherein said method is used to treat or prevent a cardiovascular disease selected from restenosis, cardiomegaly,



atherosclerosis, myocardial infarction, or congestive heart failure.

31. The method according to claim 26, wherein  
5 said method is used to treat or prevent neurodegenerative disease selected from Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, cerebral ischemia or neurodegenerative disease caused by traumatic injury, glutamate neurotoxicity or  
10 hypoxia.

32. The method according to claim 26,  
comprising the additional step of administering to said patient an additional therapeutic agent selected from an  
15 anti-proliferative agent, an anti-inflammatory agent, an immunomodulatory agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating liver disease, an anti-viral agent, an agent for treating blood disorders, an agent for treating diabetes,  
20 or an agent for treating immunodeficiency disorders, wherein:

said additional therapeutic agent is appropriate for the disease being treated; and  
said additional therapeutic agent is  
25 administered together with said composition as a single dosage form or separately from said composition as part of a multiple dosage form.

33. A composition for coating an implantable  
30 device comprising a compound according to claim 1 and a carrier suitable for coating said implantable device.

34. An implantable device coated with a composition according to claim 33.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 02/13396

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D249/06 C07D249/08 C07D249/10 C07D403/04 C07D401/14  
C07D405/14 C07D417/14 C07D403/14

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 52941 A (SEARLE & CO ;HANSON GUNNAR J (US); LIAO SHUYUAN (US)) 26 November 1998 (1998-11-26) claims	1,22-34
A	WO 95 31451 A (SMITHKLINE BEECHAM CORP ;ADAMS JERRY LEROY (US); GALLAGHER TIMOTHY) 23 November 1995 (1995-11-23) cited in the application claims	1,22-34
P,A	WO 01 56993 A (JANETKA JAMES ;VERTEX PHARMA (US); CAO JINGRONG (US); HALE MICHAEL) 9 August 2001 (2001-08-09) claims	1,22-34

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

20 June 2002

Date of mailing of the international search report

28/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 02/13396

### Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 24-32 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

#### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No  
PCT/US 02/13396

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9852941	A	26-11-1998	AU 7726898 A	11-12-1998
			EP 1019394 A1	19-07-2000
			JP 2002502380 T	22-01-2002
			US 6087381 A	11-07-2000
			WO 9852941 A1	26-11-1998
<hr/>				
WO 9531451	A	23-11-1995	US 5559137 A	24-09-1996
			EP 0871622 A1	21-10-1998
			JP 10500413 T	13-01-1998
			WO 9531451 A1	23-11-1995
			US 5998425 A	07-12-1999
			US 6306883 B1	23-10-2001
<hr/>				
WO 0156993	A	09-08-2001	AU 3488801 A	14-08-2001
			AU 3672301 A	14-08-2001
			BR 0104424 A	08-01-2002
			CZ 20013540 A3	13-03-2002
			EP 1196390 A2	17-04-2002
			EP 1200422 A2	02-05-2002
			NO 20014837 A	04-12-2001
			WO 0156993 A2	09-08-2001
			WO 0157022 A2	09-08-2001
<hr/>				